

Partial Derivatives

For a function of two independent variables, $z = f(x, y)$, the partial derivative of f or z with respect to x represented by $f_x, z_x, \partial f / \partial x$ or $\partial z / \partial x$, we can be found f_x by applying all the usual rules of differentiation. The only exception is that, whenever and wherever the second variable y appears, it is treated as a constant in every respect. The partial derivative of f or z with respect to y represented by $f_y, z_y, \partial f / \partial y$ or $\partial z / \partial y$, we can similarly be found by treating x as a constant whenever it appears.

For a function of more than two independent variables, the process of finding the partial derivative of a function is called partial differentiation. In this process, the partial derivative of a function with respect to one variable is found by keeping the other variable constant.

Example 1: Find f_x and f_y for the function

1. $f(x, y) = x^2 y^4 \quad \Leftrightarrow \quad f_x = 2xy^4 \quad \text{and} \quad f_y = 4x^2 y^3$
2. $f(x, y) = x^3 + y^2 \quad \Leftrightarrow \quad f_x = 3x^2 \quad \text{and} \quad f_y = 2y$
3. $f(x, y) = e^{2y+3} \sin 3x \quad \Leftrightarrow \quad f_x = 3e^{2y+3} \cos 3x \quad \text{and} \quad f_y = 2e^{2y+3} \sin 3x$

Example 2: Find w_x and w_y for $w(x, y) = x^2 \cos(xy)$

$$w_x = x^2(-\sin(xy)) \times y + 2x \cos(xy) = -x^2 y \sin(xy) + 2x \cos(xy)$$
$$w_y = x^2(-\sin(xy)) \times x = -x^3 \sin(xy)$$

Example 3: Find h_s and h_t for $h(s, t) = t \ln(4s^2 + 1) + t^2 \tan^{-1}(2s)$

$$h_s = \frac{8st}{4s^2 + 1} + \frac{2t^2}{1 + 4s^2} = \frac{8st + 2t^2}{4s^2 + 1}$$
$$h_t = \ln(4s^2 + 1) + 2t \tan^{-1}(2s)$$

Example 4: If $f(x, y) = \frac{x - y}{x + y}$, then show that $xf_x + yf_y = 0$

$$f_x = \frac{(x + y) - (x - y)}{(x + y)^2} = \frac{2y}{(x + y)^2}$$

$$f_y = \frac{(x+y) \times (-1) - (x-y)}{(x+y)^2} = \frac{-2x}{(x+y)^2}$$

$$xf_x + yf_y = \frac{2xy}{(x+y)^2} + \frac{-2xy}{(x+y)^2} = 0$$

Example 5: If $w = x \sin(yz) + xe^{yz}$, then show that 1. $xw_x = w$ 2. $yw_y = zw_z$

$$1. w_x = \sin(yz) + e^{yz} \quad \Leftrightarrow \quad xw_x = x \sin(yz) + xe^{yz} = w$$

$$2. w_y = xz \cos(yz) + xze^{yz} \quad \Leftrightarrow \quad yw_y = xyz \cos(yz) + xyze^{yz}$$

$$w_z = xy \cos(yz) + xye^{yz} \quad \Leftrightarrow \quad zw_z = xyz \cos(yz) + xyze^{yz}$$

$$\text{So, } yw_y = zw_z$$

Second Order Partial Derivatives

Let $z = f(x, y)$ be a function of x and y , then the second partial derivative of f with respect to x is f_{xx} , the second partial derivative of f with respect to y is f_{yy} , the second partial derivative of f with respect to y and then with respect to x is f_{xy} and the second partial derivative of f with respect to x and then with respect to y is f_{yx} .

$$\left(f_{xx} \equiv \frac{\partial^2 f}{\partial x^2}, f_{yy} \equiv \frac{\partial^2 f}{\partial y^2}, f_{xy} \equiv \frac{\partial^2 f}{\partial x \partial y} \text{ and } f_{yx} \equiv \frac{\partial^2 f}{\partial y \partial x} \right)$$

f_{xy} and f_{yx} are called mixed partial derivatives where $f_{xy} = f_{yx}$.

Example 6: Find f_{xx}, f_{xy}, f_{yx} and f_{yy} for $f(x, y) = x^2 + 3xy - y^4$

$$f_x = 2x + 3y \quad \Leftrightarrow \quad f_{xx} = 2 \quad \text{and} \quad f_{yx} = 3$$

$$f_y = 3x - 4y^3 \quad \Leftrightarrow \quad f_{yy} = -12y^2 \quad \text{and} \quad f_{xy} = 3$$

Example 7: Find f_{rr} and $f_{\theta\theta}$ for $f(r, \theta) = r^2 \sin^2 \theta$

$$f_r = 2r \sin^2 \theta \quad \Leftrightarrow \quad f_{rr} = 2 \sin^2 \theta$$

$$f_\theta = 2r^2 \sin \theta \cos \theta \quad \text{but} \quad (2 \sin \theta \cos \theta = \sin 2\theta)$$

$$f_\theta = r^2 \sin 2\theta \quad \Leftrightarrow \quad f_{\theta\theta} = 2r^2 \cos 2\theta$$

Example 8: Find all first and second order partial derivatives of the function

$$f(x, y, z) = 3x^2 - 2xy^2 + 4x^2z + z^3y + 5$$

$$f_x = 6x - 2y^2 + 8xz \quad \Leftrightarrow \quad f_{xx} = 6 + 8z, \quad f_{yx} = -4y \quad \text{and} \quad f_{zx} = 8x$$

$$f_y = -4xy + z^3 \quad \Leftrightarrow \quad f_{yy} = -4x, \quad f_{xy} = -4y \quad \text{and} \quad f_{zy} = 2z^2$$

$$f_z = 4x^2 + 2z^2y \quad \Leftrightarrow \quad f_{zz} = 4zy, \quad f_{xz} = 8x \quad \text{and} \quad f_{yz} = 2z^2$$

Chain Rule for Partial Derivatives

If $z = f(x, y)$ and $x = x(u, v)$, $y = y(u, v)$, then $z = z(u, v)$ and

$$\boxed{\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x} \times \frac{\partial x}{\partial u} + \frac{\partial z}{\partial y} \times \frac{\partial y}{\partial u}} \quad \text{and} \quad \boxed{\frac{\partial z}{\partial v} = \frac{\partial z}{\partial x} \times \frac{\partial x}{\partial v} + \frac{\partial z}{\partial y} \times \frac{\partial y}{\partial v}}$$

Example 9: If $z = x^2 + y^2$, $x = 2u + v$, $y = 2v - u$.

Find $\frac{\partial z}{\partial u}$ and $\frac{\partial z}{\partial v}$ as a functions of u and v

$$\frac{\partial z}{\partial x} = 2x = 4u + 2v, \quad \frac{\partial z}{\partial y} = 2y = 4v - 2u$$

$$\frac{\partial x}{\partial u} = 2, \quad \frac{\partial y}{\partial u} = -1, \quad \frac{\partial x}{\partial v} = 1 \quad \text{and} \quad \frac{\partial y}{\partial v} = 2$$

$$\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x} \times \frac{\partial x}{\partial u} + \frac{\partial z}{\partial y} \times \frac{\partial y}{\partial u}$$

$$= (4u + 2v) \times 2 + (4v - 2u) \times (-1)$$

$$= 8u + 4v - 4v + 2u = 10u$$

$$\frac{\partial z}{\partial v} = \frac{\partial z}{\partial x} \times \frac{\partial x}{\partial v} + \frac{\partial z}{\partial y} \times \frac{\partial y}{\partial v}$$

$$= (4u + 2v) \times 1 + (4v - 2u) \times 2$$

$$= 4u + 2v + 8v - 4u = 10v$$

Example 10: Let $w = xy - z$, $x = \sin t$, $y = \cos t$ and $z = t$. Find $\frac{\partial w}{\partial t}$

$$\frac{\partial w}{\partial t} = \frac{\partial w}{\partial x} \times \frac{\partial x}{\partial t} + \frac{\partial w}{\partial y} \times \frac{\partial y}{\partial t} + \frac{\partial w}{\partial z} \times \frac{\partial z}{\partial t}$$

$$= y \times \cos t + x \times (-\sin t) + (-1) \times 1$$

$$= \cos^2 t - \sin^2 t - 1$$

Laplace's Equation: We say that the function $f(x, y)$ satisfies Laplace's equation if

$$\boxed{\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0} \quad (f_{xx} + f_{yy} = 0)$$

Example 11: Show that the functions satisfy Laplace's equation

1. $f(x, y) = e^{-2y} \cos 2x$

2. $w(s, t) = \ln(t^2 + s^2)$

$$1. \quad \frac{\partial f}{\partial x} = -2e^{-2y} \sin 2x \quad \Leftrightarrow \quad \frac{\partial^2 f}{\partial x^2} = -4e^{-2y} \cos 2x$$

$$\frac{\partial f}{\partial y} = -2e^{-2y} \cos 2x \quad \Leftrightarrow \quad \frac{\partial^2 f}{\partial y^2} = 4e^{-2y} \cos 2x$$

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = -4e^{-2y} \cos 2x + 4e^{-2y} \cos 2x = 0$$

$$2. \quad \frac{\partial w}{\partial s} = \frac{2s}{t^2 + s^2} \quad \Leftrightarrow \quad \frac{\partial^2 w}{\partial s^2} = \frac{2(t^2 + s^2) - 2s \times 2s}{(t^2 + s^2)^2}$$

$$\frac{\partial^2 w}{\partial s^2} = \frac{2t^2 + 2s^2 - 4s^2}{(t^2 + s^2)^2} = \frac{2t^2 - 2s^2}{(t^2 + s^2)^2}$$

$$\frac{\partial w}{\partial t} = \frac{2t}{t^2 + s^2} \quad \Leftrightarrow \quad \frac{\partial^2 w}{\partial t^2} = \frac{2(t^2 + s^2) - 2t \times 2t}{(t^2 + s^2)^2}$$

$$\frac{\partial^2 w}{\partial t^2} = \frac{2t^2 + 2s^2 - 4t^2}{(t^2 + s^2)^2} = \frac{2s^2 - 2t^2}{(t^2 + s^2)^2}$$

$$\frac{\partial^2 w}{\partial s^2} + \frac{\partial^2 w}{\partial t^2} = \frac{2t^2 - 2s^2}{(t^2 + s^2)^2} + \frac{2s^2 - 2t^2}{(t^2 + s^2)^2} = 0$$

The 1-D Heat Equation: The 1-D Heat equation takes the form:

$$\boxed{\frac{\partial T}{\partial t} = k \frac{\partial^2 T}{\partial x^2}}, \text{ where } k > 0 \text{ which is called the thermal diffusivity.}$$

Example 12: Show that the function $T(x, t) = 3e^{-4\pi^2 t} \cos(2\pi x)$ satisfy heat equation, with $k = 1$.

$$\frac{\partial T}{\partial t} = -12\pi^2 e^{-4\pi^2 t} \cos(2\pi x)$$

$$\frac{\partial T}{\partial x} = -6\pi e^{-4\pi^2 t} \sin(2\pi x) \quad \Leftrightarrow \quad \frac{\partial^2 T}{\partial x^2} = -12\pi^2 e^{-4\pi^2 t} \cos(2\pi x)$$

$$\therefore \frac{\partial T}{\partial t} = \frac{\partial^2 T}{\partial x^2}$$

Example 13: If $T(x, t) = 2e^{-12t} \sin 2x$ satisfy heat equation, then find the thermal diffusivity k .

$$\frac{\partial T}{\partial t} = -24e^{-12t} \sin 2x$$

$$\frac{\partial T}{\partial x} = 4e^{-12t} \cos 2x \quad \Leftrightarrow \quad \frac{\partial^2 T}{\partial x^2} = -8e^{-12t} \sin 2x$$

$$\frac{\partial T}{\partial t} = k \frac{\partial^2 T}{\partial x^2} \quad \Leftrightarrow \quad -24e^{-12t} \sin 2x = -8ke^{-12t} \sin 2x$$

$$\therefore k = 3$$

Wave Equation: The wave equation takes the form: $u_{tt} = c^2 u_{xx}$

Example 14: Show that the function $u(x, t) = \cos(x + 2t) - \cos(x - 2t)$ satisfy the wave equation $u_{tt} = 4u_{xx}$

$$u_t = -2 \sin(x + 2t) - 2 \sin(x - 2t)$$

$$u_{tt} = -4 \cos(x + 2t) + 4 \cos(x - 2t)$$

$$u_x = -\sin(x + 2t) - \sin(x - 2t)$$

$$u_{xx} = -\cos(x + 2t) + \cos(x - 2t)$$

$$\therefore u_{tt} = 4u_{xx}$$

Exercises

1. If $w = \cos(x + y) + \sin(x - y)$ then show that $\frac{\partial^2 w}{\partial x^2} = \frac{\partial^2 w}{\partial y^2}$.

2. Find $\frac{\partial^2 z}{\partial x \partial y}$ if $z = x^2 \sin(2x - 3y)$.

3. If $z = \ln(xy)$ and $x = r \sin \theta$, $y = r \cos \theta$,

then show that $\frac{\partial z}{\partial \theta} = 2(\csc 2\theta - \cot 2\theta)$

4. If $w = (x^2 + y^2 + z^2)(y^2 + z^2)$, then find $\frac{\partial w}{\partial x}$ at $x = 1, y = 2$ and $z = 3$.

5. Show that the functions satisfy Laplace's equation

$$a) f(x, y) = e^{3x} \sin 3y \quad b) f(x, y) = x^3 - 3xy^2$$

6. If $T(x, t) = 5e^{-32\pi^2 t} \sin(4\pi x)$ satisfy heat equation, then find the thermal diffusivity k .

7. Show that the function $u(x, t) = e^{x+ct} - e^{x-ct}$ satisfy wave equation.