

Research Article

Pathogenic and Antimicrobial Properties of Aquatic Extracts of *Viscus album* and *Apium graveolens*

Enass Najem Oubaid¹, Zainab Adil Ghani Chabuck², Rafal J. Al-Saigh³, Nada Khazal Kadhim Hindi⁴ and Samah Ahmed Kadhum³

¹College of Pharmacy, University of Babylon, Babylon, Iraq

²Department of Medical Microbiology, College of Medicine, University of Babylon, Babylon, Iraq

³Department of Clinical Laboratory Sciences, College of Pharmacy, University of Babylon, Babylon, Iraq

⁴Department of Basic and Medical Science, College of Nursing, University of Babylon, Babylon, Iraq

Abstract

Background and Objective: Wild plants had been recognized to have chief roles for many populations to provide them with many requirements, also, these plants have been applied for the treatments of various complaints, especially microbial infections. This study aimed to identify the inhibitory effects of watery extracts obtained from *Viscus album* and *Apium graveolens* against the number of clinical bacterial isolates. **Materials and Methods:** Aqueous extracts of *Viscus album* and *Apium graveolens* were applied to examine their effects on inhibition of growth and some physiological parameters such as swarming and biofilm-formation against 19 clinical bacterial isolates (Gram-positive and negative). Method of agar wells-diffusion, tissue culture plate method for biofilm inhibition test, inhibition of adherence and swarming assay were applied for estimating and evaluating the inhibitory effects. **Results:** *Viscus album* and *Apium graveolens* extracts showed significant inhibitory effects against most tested Gram-positive and negative bacterial isolates with inhibitory ranges 20-35 and 18-28 mm, respectively, also with significant strong inhibition of adhesion and biofilm formation by *Viscus album* when compared with less inhibition by *Apium graveolens*. **Conclusion:** Aquatic extracts of *Viscus album* and *Apium graveolens* can be considered beneficial prophylaxis and treated against a wide range of pathogenic bacteria.

Key words: Pathogenic bacteria, antimicrobial, aquatic extract, plant extracts, *Viscus album*, *Apium graveolens*, tissue culture

Citation: Oubaid, E.N., Z.A.G. Chabuck, R.J. Al-Saigh, N.K.K. Hindi and S.A. Kadhum, 2022. Pathogenic and antimicrobial properties of aquatic extracts of *Viscus album* and *Apium graveolens*. Asian J. Plant Sci., 21: 360-68.

Corresponding Author: Rafal J. Al-Saigh, Department of Clinical Laboratory Sciences, College of Pharmacy, University of Babylon, Babylon, Iraq
Tel: +9647732258232

Copyright: © 2022 Enass Najem Oubaid et al. This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

The rapid bacterial spreading and uncontrolled usage of antibacterial agents have led to the progression of horrible drug resistance among variable virulent pathogenic bacterial species. This is resulting in the development of a huge problem of antimicrobial resistance which increases the rate of morbidity and mortality. Additionally, other problems regarding the cost and availability of different synthetic broad spectrum antibiotics, as most of them are too costly and difficult to be got. These drawbacks abate the therapeutic convenience of available synthetic antibacterial drugs and enhance the requirement to replace them with effective, available and cheaper alternative remedies¹.

Various plants can be functional for the treatments and solving of different illnesses and problems, by serving as sources of many powerful drugs. These plants contain different bio-active composites which are responsible for the therapeutic activities of these plants with their availability and diversity of them in different ecological regions around the world. Thus, many new synthetic drugs are to be obtained from natural resources, remarkably of plant origin². The antibacterial properties of aromatic and spices herbs have been detected in several types of research, as their composition, activities, applications, role as a natural food preservative and as medical treatments.

Apium graveolens belonging to the family Umbelliferae (English name Celery), is a herbaceous, biennial and branched stem medicinal plant widely available in various countries, it is ordinarily cultured in the Europe area as a diet yield and it is also grown in Egyptian cities, regions of Kingdom of Saudi Arabia and Algeria³. Celery leaves contain flavonoids, tannins and saponins which have antibacterial effects, as the juice of its roots and leaves possess effective biochemical actions⁴. This extract has many medical activities like anticancer, anti-inflammatory, anti-bacterial, anti-spasmodic, antihypercholesterolemic, anti-hepatotoxic and analgesic⁵ with many types of research that record anti-inflammatory, antibacterial and antifungal effects of essential oil of celery^{6,7}.

Celery can be a possible application to enhance the development and production of commercially used insect repellents and to be applied as alternatives for the commonly used synthetic chemicals in the community for controlling insects mainly the vector ones. In past years, a certain substance called CAH had been detected in the ethanolic extract of celery seed, which had been checked as an active reagent for *Helicobacter pylori* infection treatment⁸. Additionally, Celery has powerful antimicrobial activities against *Saccharomyces cerevisiae*, *Bacillus subtilis* and *E. coli*, the oil of celery seeds was detected to be effective against

Campylobacter jejuni, with potent inhibitory effects against various saprophytic and pathogenic microorganisms^{9,10}. Moreover, celery contains two senkyunolides and sedanolide which are effective against mosquito larvae, nematodes and fungi, furanocoumarins inhibited *Micrococcus luteus*, *Listeria monocytogenes* and *E. coli* O157:H7^{11,12}.

Viscum album (Mistletoe) is a member of the Loranthaceae family. It is a familiar evergreen parasitic plant, growing on deciduous trees as a ball-like scrub. The evergreen plants are considered obligate parasites, depending partly on their host to get minerals and water but can process photosynthesis. Thus it is considered a semi-parasite that can live in most regions of the world, either on edible or nonedible trees, remembering that only those that grow on edible plants can be used for medicinal purposes¹³.

It contains resin, saponin, organic acids and alkaloids and also contains hydrophilic cytotoxic substances and immunomodulatory proteins such as viscotoxins and mistletoe lectins. They have the ability of immune system stimulation via leukocytes activation leading to the release of cytokines, inhibiting cellular proliferation with in vitro and in vivo apoptosis induction^{14,15}.

The certain study applied in vivo on rats infected by *S. aureus* with *B. cereus* and *E. coli* with *P. aeruginosa*. Haematological, bacterial and histopathological studies, 7 days after treatment with extract of *Viscum album* exhibited its therapeutic effect¹.

This work was done to examine the antibacterial inhibitory activity of aquatic extract of both *Viscus Album* and *Apium graveolens* against a group of clinically isolated bacteria, whether by direct inhibition or inhibition of adhesion and biofilm formation.

MATERIALS AND METHODS

Plant collection: Fresh dried plant materials of *Viscus album* (mistletoe) and *Apium graveolens* (celery) were purchased from a local market in Hilla city/ Iraq. Preparation of aquatic extract from fresh *Viscus album* and *Apium graveolens* was prepared by soaking 50 gm of each one as a dried powder in 100 mL distilled water and allowed to stand for about 72 hrs then sterilized by filtration (by Millipore filter paper-0.45). The above-prepared extracts were cleaned, peeled, heat dried and cut into small pieces, the electric blender was used to make the ground and placed in clean containers¹⁶.

Bacterial and isolates: Nineteen bacterial isolates obtained from clinical specimens (Gram +ve and Gram -ve isolates)

Table 1: Gram-positive and Gram-negative bacteria involved in this study

Gram-negative bacteria	Gram-positive bacteria
<i>Salmonella typhi</i>	<i>Staphylococcus aureus</i>
<i>Salmonellatyphimurum</i>	<i>Staphylococcus epidermidis</i>
<i>Pseudomonas aeruginosa</i>	<i>Staphylococcus saprophyticus</i>
<i>Pseudomonas fluorescences</i>	<i>Streptococcus pyogenes</i>
<i>Proteus vulgaris</i>	<i>Streptococcus pneumoniae</i>
<i>Proteus mirabilis</i>	<i>Streptococcus mutans</i>
<i>Klebsiella pneumoniae</i>	<i>Streptococcus faecalis</i>
<i>Enterobacter aerogenes</i>	<i>Streptococcus agalactiae</i>
<i>Acinetobacter</i>	
<i>Escherichia coli Serratia spp.</i>	

Table 2: Bacterial adherence and biofilm formation by the method of TCP (20)

Mean of OD value at 630 nm	Adherence	Biofilm formation
0.120>	Non	Non
0.120-0.240	Moderate	Moderate
>0.240	Strong	High

were used in the work (Table 1). Bacterial isolates were activated three continual times on nutrient agar and then stored at 4EC as nutrient agar slant. The documentation of isolates was established by various biochemical tests (Api system (Biomeraux, France) in microbiological laboratory/college of Pharmacy/the University of Babylon¹⁷.

Antimicrobial activity test by agar-well diffusion assay (in vitro)

Antibacterial activity assay: According to CLSI¹⁷, the screening of antimicrobial activities of each aquatic extract on the tested bacteria was determined on Muller Hinton agar media by using the agar well diffusion method. Wells of 8 mm diameter were made on the solid agar using a sterile glass borer. Approximately 20 µL of each extract was inoculated onto wells and were made in the spread plate culture of each microbial isolate, then incubated at 37EC for 24 hrs. The examination of the diameter of the inhibitory zone by measuring it in millimetres and comparing them with the inhibitory zone of Ciprofloxacin (2 µg mL⁻¹) (the test was performed in triplicates)^{16,18}.

Biofilm formation assay: Semi-quantitative microtiter plate test or tissue culture plate method assay (TCP) designated by previous studies¹⁹ was assumed as the gold standard method for detection of biofilm formation²⁰. Results can be compared in Table 2.

Adherence test: Bacterial adhesion to the epithelial cell of the mouth is one of the chief and important virulence criteria of those bacteria and can be detected using the method suggested by authors^{21,22}.

Statistical analysis: To analyze the data, Bonferroni test²³ was applied, ($p \leq 0.05$) to show significant differences between the types of extracts and with ciprofloxacin by one-way ANOVA. All statistical analysis was performed with the Software Prism 5.01 (GraphPad Inc., La Jolla, CA).

RESULTS

The results of the two aquatic extracts were compared with the Ciprofloxacin (2 µg mL⁻¹) against the same Gram positive and negative bacteria and it showed less inhibitory effect against most tested bacteria than those two aquatic extracts (Fig. 1a-k and Fig. 2a-h). The two aquatic extracts showed a significant antibacterial effect. *Viscus album* observed high efficacy against both Gram-negative and Gram-positive bacteria with an inhibition zone ranging from 20 mm when used against *Pseudomonas aeruginosa* and *Pseudomonas fluorescences* (Fig. 1c and Fig. 1e, respectively) to 35 mm on use against *Streptococcus agalactia* (Fig. 2h). While the antibacterial effect of *Apium graveolens* revealed a slight antibacterial effect against both Gram-negative and positive bacteria at a range of 18 for *Pseudomonas aeruginosa* (Fig. 1c) to 28 mm for *Streptococcus faecalis* (Fig. 2g).

Regarding the detection of anti-adherence and antibiofilm activities of aquatic extracts of *Viscus album* against Gram-negative bacteria, it was showing strong inhibition to the growth with the adhesion of bacterial isolate on a glass plate (Table 3), with strong inhibition of adhesion of epithelial cell for all using bacteria (Table 4). In contrast to moderate inhibition for both anti-adherence and anti-biofilm activities by using *Apium graveolens* extract.




Fig. 1(a-k): Antibacterial effect of *Viscus album* and *Apium graveolens* against several clinical isolates of Gram-negative bacteria by agar well diffusion method in comparison with Ciprofloxacin (2 µg mL⁻¹), (a) *Salmonella typhi*, (b) *Salmonella typhimurium*, (c) *Pseudomonas aeruginosa*, (d) *Escherichia coli*, (e) *Pseudomonas fluorescens*, (f) *Proteus vulgaris*, (g) *Proteus mirabilis*, (h) *Serratia*, (i) *Klebsiella pneumoniae*, (j) *Enterobacter aerogenes* and (k) *Acinetobacter*.

Number of asterisks (*) corresponds to the level of the statistical significance (*p<0.05, **p<0.01 and ***p<0.001)

Table 3: Anti-adherence activity of aquatic extracts of *Viscus album* and *Apium graveolens* against Gram-negative bacteria

Gram-negative bacteria	<i>Viscus album</i>	<i>Apium graveolens</i>
<i>Salmonella typhi</i>	High	Moderate
<i>Salmonella typhimurium</i>	High	Moderate
<i>Pseudomonas aeruginosa</i>	High	Moderate
<i>Pseudomonas fluorescens</i>	High	Moderate
<i>Proteus vulgaris</i>	High	Moderate
<i>Proteus mirabilis</i>	High	Moderate
<i>Klebsiella pneumoniae</i>	High	Moderate
<i>Enterobacter aerogenes</i>	High	Moderate
<i>Acinetobacter</i>	High	Moderate
<i>Escherichia coli</i>	High	Moderate
<i>Serratia spp.</i>	High	Moderate




Fig. 2(a-h): Antibacterial effect of *Viscus album* and *Apium graveolens* against the number of a clinical isolate of Gram-positive bacteria by agar well method in comparison with Ciprofloxacin (2 µg mL⁻¹), (a) *Staphylococcus aureus*, (b) *Staphylococcus epidermidis*, (c) *Staphylococcus saprophyticus*, (d) *Streptococcus pyogenes*, (e) *Streptococcus pneumoniae*, (f) *Streptococcus mutans*, (g) *Streptococcus faecalis* and (h) *Streptococcus agalactiae*. Number of asterisks (*) corresponds to the level of the statistical significance (*p<0.05, **p<0.01 and ***p<0.001)

Table 4: Anti-biofilm activity of aquatic extracts of *Viscus album* and *Apium graveolens* against Gram-negative bacteria

Gram-negative bacteria	<i>Viscus album</i>	<i>Apium graveolens</i>
Salmonella typhi	High	Moderate
Salmonella typhimurium	High	Moderate
Pseudomonas aeruginosa	High	Moderate
Pseudomonas fluorescences	High	Moderate
Proteus vulgaris	High	Moderate
Proteus mirabilis	High	Moderate
Klebsiella pneumoniae	High	Moderate
Enterobacter aerogenes	High	Moderate
Acinetobacter	High	Moderate
Escherichia coli	High	Moderate
Serratia spp.	High	Moderate

DISCUSSION

Medicinally-used plants in particular display a prolonged history in the management of different diseases with a foremost role in the development and production of novel drugs today. While medicine and technology have been developed widely, with an

increment in poverty, decrement in natural richness and other obstacles, all of these enhance the obligatory usage of natural products for many goals.

Hildegard von Bingen explained the use of mistletoe as a treatment for splenic and hepatic diseases. Mistletoe was also used for the cure of mumps, leprosy, hepatitis, affections of the liver and lungs, deworming of children, treatment of labour pains, fractures and gout²⁴.

Many studies were carried out to detect its antimicrobial activities against different microbes²⁵⁻³⁰. Some researchers had shown that antibacterial activities of these extracts were more effective and obvious against most Gram-negative pathogenic bacteria because of their anti-biofilm activity when compared with Gram-positive bacterial species that have thin biofilms^{26,31,32}.

Orhue et al.³³ used different alcoholic mistletoe extracts against *Klebsiella aerogenes*, *P. aeruginosa*, *E. coli* and *S. aureus*, they revealed antibacterial activity to *P. aeruginosa* and the *Staph. aureus* only. While Oguntoye et al.³⁴ showed

that aquatic extract produced inhibition against bacterial isolates more than the ethanolic extract.

Viscus album not only inhibits bacterial growth but also its compounds stimulate the immune system via activation of granulocytes, monocytes/macrophages, T-cells, natural killer (NK) cells and dendritic cells, with induction of various cytokines such as TNF- α , GM-CSF, IFN- γ IL-1, IL-2, IL-4, IL-5, IL-6, IL-8, IL-10 and IL-12^{24,35}.

These wide different pharmaceutical applications are the result of wealthy chemical structures of *Viscum* species. The chief dynamic active compounds for these species are flavonoids, lectins, Visco toxins, phenolic acids, sterols, fatty acids terpenoids, alkaloids, phenylpropanoids and lignans³⁶.

Regarding *Apium graveolens*, Genatrika et al.⁴ tried to utilize its extract as toothpaste to inhibit *Streptococcus mutans* and it showed significant antibacterial activity against these bacteria, moreover, they showed that it has a protective or disease preventive antibacterial activities. Furthermore, Venskutonis et al.³⁷ showed that celery root and leaf extracts have inhibited the growth of bacteria in vitro, moreover, Gram-negative bacteria were the commonest bacteria affected by these extracts with different concentrations.

Certain studies have shown that celery improves digestion, cardiovascular disorders, central nervous system, with additional anti-inflammatory, antimicrobial and other various pharmacological activities^{8,38-40}. Moreover, certain work studied the effects of this plant extract on *M. tuberculosis* by the use of 200 mg mL⁻¹ of whole-plant extract made using 70% ethanol, it showed inhibitory effects on MTB (only 20 colonies appeared after an incubation period of 4 weeks)⁴⁰.

Flavonoids, Saponins and Alkaloids components of celery and their role as antibacterial effects were also studied⁴¹. Its phytochemicals were detected in the leaves of celery⁴². Flavonoids exhibit antibacterial effects via different mechanisms such as energy metabolism, inhibition of cytoplasmic membrane function and inhibition of nucleic acid synthesis⁴³. Saponins' antibacterial effects seem to have membranolytic properties⁴⁴, as it has increased the bacterial cell membrane permeability without destroying them⁴⁵. Moreover, the mechanism of the action of highly aromatic alkaloids can intercalate with DNA⁴⁶. Both above mechanisms may explain the suggested mechanisms of antibacterial effectivity of our plant's extracts.

CONCLUSION

The infections caused by highly resistant Gram-positive and negative bacteria are among the most difficult diseases all over the

world. Medicinal plants that can be used traditionally to treat such infections are suitable to face these problems.

SIGNIFICANCE STATEMENT

Therefore, aquatic extracts of *Viscus Album* and *Apium graveolens* can be considered beneficial for prophylaxis and treatment against a wide range of pathogenic bacteria. Moreover, these plant extracts can be a source of new complementary medicines and support for the treatment of many infectious diseases and can prevent or decrease their complications.

REFERENCES

1. Szurpnicka, A., A. Kowalczyk and A. Szerk, 2020. Biological activity of mistletoe: In vitro and in vivo studies and mechanisms of action. Arch. Pharm. Res., 43: 593-629.
2. Uddin, Z., A.A. Shad, J. Bakht, I. Ullah and S. Jan, 2015. In vitro antimicrobial, antioxidant activity and phytochemical screening of *Apium graveolens*. Pak. J. Pharm. Sci., 28: 1699-1704.
3. Al-Asmari, A.K., A.M. Al-Elaiwi, M.T. Athar, M. Tariq, A.A. Eid and S.M. Al-Asmary, 2014. A review of hepatoprotective plants used in Saudi traditional medicine. Evidence-Based Complementary Altern. Med., Vol. 2014. 10.1155/2014/890842.
4. Genatrika, E., F. Satriani and I. Hapsari, 2019. Antibacterial activity of celery leaves (*Apium graveolens* L.) formulated in toothpaste against *Streptococcus mutans*. Int. J. App. Pharm., 11: 14-16.
5. Modaresi, M., G. Ghalamkari and A. Jalalizand, 2012. The effect of celery (*Apium graveolens*) extract on the reproductive hormones in male mice. APCBEE Procedia, 4: 99-104.
6. Mencherini, T., A. Cau, G. Bianco, R.D. Loggia, R.P. Aquino and G. Autore, 2007. An extract of *Apium graveolens* var. dulce leaves: Structure of the major constituent, apiin and its antiinflammatory properties. J. Pharm. Pharmacol., 59: 891-897.
7. Edziri, H., S. Ammar, L. Souad, M.A. Mahjoub and M. Mastouri et al., 2012. In vitro evaluation of antimicrobial and antioxidant activities of some Tunisian vegetables. S. Afr. J. Bot., 78: 252-256.
8. Al-Asmari, A., M.T. Athar and S.G. Kadasah, 2017. An updated phytopharmacological review on medicinal plant of Arab Region: *Apium graveolens* Linn. Phcog. Rev., 11: 13-18.
9. Elgayyar, M., F.A. Draughon, D.A. Golden and J.R. Mount, 2001. Antimicrobial activity of essential oils from plants against selected pathogenic and saprophytic microorganisms. J. Food Prot., 64: 1019-1024.
10. Friedman, M., P.R. Henika and R.E. Mandrell, 2002. Bactericidal activities of plant essential oils and some of their isolated constituents against *Campylobacter jejuni*, *Escherichia coli*,

- Listeria monocytogenes* and *Salmonella enterica*. J. Food Prot., 65: 1545-1560.
11. Momin, R.A. and G. Nair, 2001. Mosquitocidal, nematicidal, and antifungal compounds from *Apium graveolens* L. seeds. J. Agric. Food Chem., 49: 142-145.
 12. Shad, A.A., H.U. Shah, J. Bakht, M.I. Choudhary and J. Ullah, 2011. Nutraceutical potential and bioassay of *Apium graveolens* L. grown in Khyber Pakhtunkhwa-Pakistan. J. Med. Plants Res., 5: 5160-5166.
 13. Evans, J., 2005. Mistletoe: Good for more than free kisses. J. Am. Bot. Council., 68: 50-59.
 14. Urech, K., G. Schaller and C. Jäggy, 2006. Viscotoxins, mistletoe lectins and their isoforms in mistletoe (*Viscum album* L.) extracts iscador. *Arzneim.-Forsch.*, 56: 428-434.
 15. Hajtó, T., K. Fodor, P. Perjési and P. Németh, 2011. Difficulties and perspectives of immunomodulatory therapy with mistletoe lectins and standardized mistletoe extracts in evidence-based medicine. *Evidence-Based Complementary Altern. Med.*, Vol. 2011. 10.1093/ecam/nep191.
 16. Chabuck, Z., B. Hadi and N. Hindi, 2018. Evaluation of antimicrobial activity of different aquatic extracts against bacterial isolates from UTI in Babylon Province, Iraq. *J. Pure Appl. Microbiol.*, 12: 693-700.
 17. Forbes, B.A., D.F. Sahn, W.R. Bailey, A.S. Weissfeld and E.G. Scott, 2007. *Bailey and Scotts Diagnostic Microbiology*. 12th Edn., Elsevier Mosby, Missouri, US, ISBN: 9780323030656, Pages: 1031.
 18. Hindi, N.K.K. and Z.A.G. Chabuck, 2013. Antimicrobial activity of different aqueous lemon extracts. *J. Appl. Pharm. Sci.*, 3: 074-078.
 19. Christensen, G.D., W.A. Simpson, J.J. Younger, L.M. Baddour, F.F. Barrett, D.M. Melton and E.H. Beachey, 1985. Adherence of coagulase-negative staphylococci to plastic tissue culture plates: A quantitative model for the adherence of staphylococci to medical devices. *J. Clin. Microbiol.*, 22: 996-1006.
 20. Girish, V.M., H. Liang, J.T. Aguilan, J.D. Nosanchuk, J.M. Friedman and P. Nacharaju, 2019. Anti-biofilm activity of garlic extract loaded nanoparticles. *Nanomed. Nanotechnol. Biol. Med.*, Vol. 20. 10.1016/j.nano.2019.04.012.
 21. Avila-Campos, M.J., M.R.L. Simionato, C.A.I. Silvana, M.P.A. Mayer, J.L. De Lorenzo, F. Zelante, 2000. Virulence factors of *Actinobacillus actinomycetemcomitans*: Other putative factors. *Pesqui. Odontol. Bras.*, 14: 5-11.
 22. Li, L., D. Matevski, M. Aspiras, R.P. Ellen and G. Lepine, 2004. Two epithelial cell invasion-related loci of the oral pathogen *Actinobacillus actinomycetemcomitans*. *Oral Microbiol. Immunol.*, 19: 16-25.
 23. Daniel, W.W., 2009. *Biostatistics: A Foundation for Analysis in the Health Sciences*. 9th Edn., John Wiley and Sons, Canada, ISBN: 9780470105825, Pages: 960.
 24. Büssing, A., 2000. *Mistletoe: The Genus Viscum*. Harwood Academic, Netherlands, ISBN: 90-5823-092-9, Pages: 263.
 25. Sengul, M., H. Yildiz, N. Gungor, B. Cetin, Z. Eser and S. Ercisli, 2009. Total phenolic content, antioxidant and antimicrobial activities of some medicinal plants. *Pak. J. Pharm. Sci.*, 22: 102-106.
 26. García-García, J.D., J.C. Anguiano-Cabello, R. ArredondoValdés, C.A.C. del Toro and J.L. Martínez-Hernández et al., 2021. Phytochemical characterization of *Phoradendron bollanum* and *Viscum album* subs. *austriacum* as Mexican mistletoe plants with antimicrobial activity. *Plants*, Vol. 10. 10.3390/plants10071299.
 27. Assaf, A.M., R.N. Haddadin, N.A. Aldouri, R. Alabbassi, S. Mashallah, M. Mohammad and Y. Bustanji, 2013. Anticancer, anti-inflammatory and anti-microbial activities of plant extracts used against hematological tumors in traditional medicine of Jordan. *J. Ethnopharmacol.*, 145: 728-736.
 28. Abualhasan, M., N. Jaradat, N. Abu-Hasan, M. Almasri and A.A. Taha et al, 2014. Bioactivity of *Viscum album* extracts from olive and almond host plants in Palestine. *Phcog. J.*, 6: 117-123.
 29. Kusi, M., K. Shrestha and R. Malla, 2015. Study on phytochemical, antibacterial, antioxidant and toxicity profile of *Viscum album* Linn associated with *Acacia catechu*. *Nepal J. Biotechnol.*, 3: 60-65.
 30. Shah, S., Y. Rehman, A. Iqbal, Z. Rahman, B. Zhou, M. Peng and Z. Li, 2017. Phytochemical screening and antimicrobial activities of stem, leaves and fruit extracts of *Viscum album* L. *J. Pure Appl. Microbiol.*, 11: 1337-1349.
 31. Erturk, Ö., H. Kati, N. Yayli and Z. Demirbag, 2003. Antimicrobial activity of *Viscum album* L. subsp. *abietis* (Wiesb). *Turk. J. Biol.*, 27: 255-258.
 32. Singh, V.K., A. Mishra and B. Jha, 2017. Anti-quorum sensing and anti-biofilm activity of *Delftia tsuruhatensis* extract by attenuating the quorum sensing-controlled virulence factor production in *Pseudomonas aeruginosa*. *Front. Cell. Infect. Microbiol.*, Vol. 7. 10.3389/fcimb.2017.00337.
 33. Orhue, P.O., E.C. Edomwande, E. Igbiosa, A.R. Momoh and O. Asekomhe, 2014. Antibacterial activity of extracts of mistletoe (*Tapinanthus dodoneifollus* (Dc) Dancer) from cocoa tree (*Theobroma cacao*). *Int. J. Herbs Pharmacol. Res.*, 3: 24-29.
 34. Oguntoye, S.O., G.A. Olatunji, O.M. Kolawole and K.I. Enonbun, 2008. Phytochemical screening and antibacterial activity of *Viscum album* (Mistletoe) extracts. *Plant Sci. Res.*, 1: 44-46.
 35. Kienle, G.S. and H. Kiene, 2003. *Mistletoe in Oncology: Facts and Conceptual Foundations*. 1st Edn., Schattauer, Stuttgart, Germany, ISBN-13: 978-3794522828, Pages: 760.
 36. Szurpnicka, A., J.K. Zjawiony and A. Sztark, 2019. Therapeutic potential of mistletoe in CNS-related neurological disorders and the chemical composition of *Viscum* species. *J. Ethnopharmacol.*, 231: 241-252.

37. Sipailiene, A., P.R. Venskutonis, A. Sarkinas and V. Cypiene, 2005. Composition and antimicrobial activity of celery (*Apium graveolens*) leaf and root extracts obtained with liquid carbon dioxide. *Acta Hortic.*, 3: 71-77.
38. Emad, A.M., D.M. Rasheed, R.F. El-Kased and D.M. El-Kersh, 2022. Antioxidant, antimicrobial activities and characterization of polyphenol-enriched extract of Egyptian celery (*Apium graveolens* L., Apiaceae) aerial parts via UPLC/ESI/TOF-MS. *Molecules*, Vol. 27. 10.3390/molecules 27030698.
39. Tashakori-Sabzevar, F., B.M. Razavi, M. Imenshahidi, M. Daneshmandi, H. Fatehi, Y.E. Sarkarizi and S.A. Mohajeri, 2016. Evaluation of mechanism for antihypertensive and vasorelaxant effects of hexanic and hydroalcoholic extracts of celery seed in normotensive and hypertensive rats. *Rev. Bras. de Farmacognosia*, 26: 619-626.
40. Kooti, W. and N. Daraei, 2017. A review of the antioxidant activity of celery (*Apium graveolens* L.). *J. Evid. Complementary Altern. Med.*, 22: 1029-1034.
41. Abbas, M.S., 2011. Study the effect of some plants extracts on growth of *Mycobacterium tuberculosis* in comparison with ethambutol. *Iraqi J. Vet. Med.*, 35: 129-134.
42. Nwaogu, L.A., C.S. Alisi, C.O. Ibegbulem and C.U. Igwe, 2007. Phytochemical and antimicrobial activity of ethanolic extract of *Landolphia owariensis* leaf. *Afr. J. Biotechnol.*, 6: 890-893.
43. Manandhar, S., S. Luitel and R.K. Dahal, 2019. In vitro antimicrobial activity of some medicinal plants against human pathogenic bacteria. *J. Trop. Med.*, Vol. 2019. 10.1155/2019/1895340.
44. Cushnie, T.P.T. and A.J. Lamb, 2005. Antimicrobial activity of flavonoids. *Int. J. Antimicrob. Agents*, 26: 343-356.
45. Al-Bayati, F.A. and H.F. Al-Mola, 2008. Antibacterial and antifungal activities of different parts of *Tribulus terrestris* L. growing in Iraq. *J. Zhejiang Univ. Sci. B*, 9: 154-159.
46. Arabski, M., A. Wągierek-Ciuk, G. Czerwonka, A. Lankoff and W. Kaca, 2012. Effects of saponins against clinical *E. coli* strains and eukaryotic cell line. *J. Biomed. Biotechnol.*, Vol. 2012. 10.1155/2012/286216.