

Prevalence of Histopathological Diagnosis of Benign and Malignant Breast Lesions in Al-Muthanna Province for Two Years Duration

Sarah Ali Abed¹; Ali Adnan Jabbar²; Roaa Salih Mahdi³; Ali Mohammed Khaleel⁴;
Suha Suhiel Hussein⁵; Maha Khadum Ghayadh⁶

¹Lecturer, College of Medicine, Al-muthanna University/Iraq, ² Ass. Prof. Dr. College of Medicine, Al-muthanna University/Iraq, ³Lecturer, College of Medicine, University of Babylon/Iraq, ⁴Dr. Specialist F.I.B.M.S., Babylon health director /MOH/Iraq, ⁵Dr. Specialist M.SC., Al- Muthanna health director/MOH/Iraq, ⁶Dr. MD (MPATH, USM, MALASIA), Al- Muthanna Health Director/MOH/Iraq

Abstract

Introduction: Breast cancer (BC) is a common form of cancer among women globally. It is considered 5th cause of death in females with equally to 522,000 patients / year. Nowadays it is considered 15.4% of death in developed countries after lung cancer. In developing nations, breast cancer considered the 1st cause of death for females 324,000 cases of deaths (14.3%) for total deaths. **Method:** One hundred and six female patients with breast lesion were collected from the Al-Hussein Teaching Hospital Laboratory/Histopathology department –Al-Muthanna province, during the period from January 2018 to January 2020. The data for cases were collected to study the age, type of breast lesion whether benign or malignant. Haematoxylin/ Eosin staining done in Al-Hussein Teaching Hospital Laboratory / Histopathology department. **Results:** Cross sectional study for 106 patients done in Al- Muthanna proven for assessment breast mass and identify malignant and benign lesion, mean age of patients was (39.4 ± 15.8) years old, with min age 13 years and max age was 81 years old. Types of biopsy taken from surgeon were 71% excisional biopsy, 21% mastectomy and 8% true cut biopsy. After pathological assessment of biopsies showed 51% malignant and 49% benign. In addition, this assessment distributed as following: fibroadenoma 31.1%, IDC/ grade II 29.2%, IDC/ grade III 9.4%, ILC and fibrocystic changes 5.7% and IDC grade I 3.8% and other types of malignant and benign after pathological assessment. Significant association between age groups and cancer of breast. **Conclusion:** After pathological assessment of biopsies, malignant breast cancer (IDC/ grade II, IDC/ grade III, ILC and fibrocystic changes) most common changes, significant association between age group and breast cancer development more age group 41- 50 years old and then (31- 40), (51– 60) years old respectively.

Key words: Prevalence, histopathological diagnosis, breast lesions, Al-muthanna province

Introduction

Breast cancer (BC) is a common form of cancer among women globally. It is considered 5th cause of death in females with equally to 522,000 patients / year ⁽¹⁾. Nowadays it is considered 15.4% of death in developed countries after lung cancer ⁽²⁾. In developing nations, breast cancer considered the 1st cause of death for females 324,000 cases of deaths (14.3%) for total

deaths ^(2, 3). This rate changes from 6 – 20 / 100.000 in West Africa and East Asia ⁽⁴⁾. The incidence in 2012 in females reach to 1.7 million (25% of entirely cancers), 883,000 patients in developed country in contradiction of 794,000 in developing nations ^(4, 5).

The incidence was increase in age after 35 years old and peaking in 60 years old ⁽⁶⁾.

Mortality rate depend on age of females, staging of malignancy, treatment respond, metastasis of malignancy, main reasons of breast cancer is hormonal factors, genetic tendencies, behaviors and ecological reasons ⁽⁷⁾.

Corresponding author:

Sarah Ali Abed

qaisajam1981@gmail.com

Benign Breast Diseases (BBDs) is not cancer. Most usual reason of breast difficulty in women and more usual than malignancy (8, 9). It is more usual than malignancy in western countries (10). More than 30% of women with BBD need treatment where survive (11). A three-ways for evaluation done by clinical assessment: US and mammogram, pathological assessment, central needle biopsy. Most women with BBD not have risk of development of breast cancer, so early diagnosis and treatment is important to remove the anxiety especially women with family history of breast cancer and must follow up annually. Most common BBD is fibro adenoma (12). Two types of breast cancer (cancer in situ): **Ductal carcinoma in situ. (DCIS)** 83% of all cases 2010-2014 unusual cells substitute usual epithelial cells around ducts of breast and metastasis to the lobules in addition to the ducts; it is can or can not developed to the aggressive malignancy it is grow slowly without any management it is misdiagnosis with benign 20%-53% diagnosed with an invasive breast malignance for 10 years or more. **Lobular carcinoma in situ.**: 13 % of patients, atypical cells developing inside and growing about some of breast lobules. It is not precursor of invasive cancer, but considered one of strong cause of invasive cancer. The aim of study is to show proportion of benign and malignant breast lesions and types of malignancy tissues by pathological assessment and relation of malignancy with increase age of females (13-15).

Method

One hundred and six female patients with breast lesion were collected from the Al-Hussein Teaching Hospital Laboratory/Histopathology department –Al-Muthanna province, during the period from January 2018 to January 2020. The data for cases were collected to study the age, type of breast lesion whether benign or malignant. Haematoxylin/Eosin staining done in Al-Hussein Teaching Hospital Laboratory / Histopathology department.

Specimens:

The one hundred and six cases classified into benign and malignant breast lesions. From each formalin fixed paraffin embedded tissue, one section of 5-micron thickness was obtained and stained by haematoxylin and eosin staining method for evaluation of morphology.

Methods of staining procedures:

The following steps were applied for (H&E) staining

method.

a) Deparaffinization: This done by adding the following:

1. 5 min. period adding Xylene.
2. 5 min. period adding Xylene.
3. 5 min. period adding 99 % ethanol
4. 5 min. period adding 99 % ethanol
5. 5 min. period adding 99 % ethanol
6. 5 min. period adding 99 % ethanol
7. 5 min. period adding 99 % ethanol
8. 5 min. period adding 95 % ethanol
9. 5 min. period adding 70 % ethanol
10. Purified water.

b) Hematoxyline and eosin staining method:

1. Dewax sections (deparaffinization as above).
2. Stain in hematoxyline for 3-10 minutes.
3. Wash well in running tap water.
4. Remove excess stain by differentiating the sections in 1% acid alcohol (1% in HCL 70% alcohol) for 5-10 seconds.
5. Wash well with in tap water until sections regain their blue color.
6. Stain in eosin for 2-5 minutes.
7. Dehydrate slowly through increasing grades of alcohol (i.e.70%, 90% and 100%).
8. Clearing by xylene.
9. Mount wit DPX.

Statistical analysis done by SPSS 22 calculated mean and SD with percentage and frequency. Chi square use for revealed association between age groups and behavior of tumor, significant association when P-value less than 0.05.

Results

Cross sectional study for 106 patients done in Al-

Muthanna proven for assessment breast mass and identify malignant and benign lesion, mean age of patients was (39.4 ± 15.8) years old, with min age 13 years and max age was 81 years old. Types of biopsy taken from surgeon were 71% excisional biopsy, 21% mastectomy and 8% true cut biopsy as showed in fig (1). After pathological assessment of biopsies showed 51% malignant and 49% benign as in fig (2). In addition, this assessment distributed as following: fibroadenoma 31.1%, IDC/ grade II 29.2%, IDC/ grade III 9.4%, ILC and fibrocystic changes 5.7% and IDC grade I 3.8% and other types of malignant and benign after pathological assessment showed in fig (3).




Fig (1): distribution of types of biopsy.




Fig (2): distribution of behavior of biopsy after pathological assessment.





Fig (3): distribution of benign and malignant according to types after pathological assessment.

According to association between ages groups and behavior of lesion either benign or malignant, there is significant association between age groups and cancer of breast, 37% of females in age group 41- 50 years old with malignant breast cancer, while 20% of malignancy in age group (31 – 40) and (51 – 60) years old. 13% of malignancy in age group 61 – 70 years old and only 6% in female over 71 years old age. As show in table (1).

Table (1): association between age groups and behavior of lesion either benign or malignant.

benign malignant		behavior			Total
Age	11 - 20 years	Count	13	0	13
		% within behavior	25.0%	0.0%	12.3%
	21 - 30 years	Count	22	2	24
		% within behavior	42.3%	3.7%	22.6%
	31 - 40 years	Count	11	11	22
		% within behavior	21.2%	20.4%	20.8%
	41 - 50 years	Count	6	20	26
		% within behavior	11.5%	37.0%	24.5%
	51 - 60 years	Count	0	11	11
		% within behavior	0.0%	20.4%	10.4%
	61 - 70 years	Count	0	7	7
		% within behavior	0.0%	13.0%	6.6%
	more than 71 years	Count	0	3	3
		% within behavior	0.0%	5.6%	2.8%
Total		Count	52	54	106
% within behavior		100.0%	100.0%	100.0%	

Pearson Chi-Square= 58.188, P-value = 0.0001** (significant).

Fig (4): Invasive ductal carcinoma with Hematoxylin & Eosin staining (10X)**Fig (5): Granulomatous mastitis with Hematoxylin & Eosin staining (4X)**

Discussion

Cancer is illnesses that lead the cells in body loss of control and change. All types of cancer cells finally changes to lump or mass that named cancer, and this malignant tumor called according to the site form it the tumor originate. Breast cancer start from gland of breast tissue that lead to production of milk called (terminal duct lobular unit). Other parts of breast tissue consist of connective, fatty and lymphatic tissues (16).

According to our study, mean age of patients was (39.4 ± 15.8) years old, with min. age 13 years and max. age was 81 years old. These results similar to study done by Augustin et al. showed the age extended since 16 - 90 years by a mean of 45.83 ± 13.5 years old (17). Zhi-Gang Yu et.al. Also, have the same results that mean age of patients was (44 ± 11.6) years old (18) characteristics and related factors of breast cancer among women in Eastern China. A total of 122,058 female subjects completed the study, with 320 confirmed cases of breast cancer (crude prevalence: 262.5/100,000; standardized prevalence: 207.7/100,000).

In our study the types of biopsy taken from surgeon were 71% excisional biopsy, 21% mastectomy and 8% true cut biopsy this is similar to Augustin et al. that showed, the total of cases that take and examine by pathologist contain of 69% of lumpectomy then 30% biopsy of breast.

After pathological assessment of biopsies in our study showed 51% malignant and 49% benign while in another study the proportion was around 4:1 or 5:1 (19), this different is due to sample collection or may be due to high no. of malignancy of breast in al-muthanna province.

In addition, this assessment distributed as following: fibroadenoma 31.1%, IDC/ grade II 29.2%, IDC/ grade III 9.4%, ILC and fibrocystic changes 5.7% and IDC grade I 3.8% and other types of malignant and benign after pathological assessment this results similar to Augustin et al. that showed he predominant tumor histological designs IDC 113 (64.9%), ILC 17 (9.8%) then histological designs were invasive ductal carcinoma 113 (64.9%), invasive lobular carcinoma 17 ductal carcinoma in situ 10 (5.7%) and medullary ca. 5 (2.8%) (17). Similar features diagnosed by Ohene et al. (20); detailed that 53.7% of grade III, 31.5% of grade II and 14.8% of grade I. Essiben et al. showed a propensity in the following course of incidence: grades II, I besides III to the Yaoundé Gynecological Obstetric and Pediatric Hospital (21) which is a referral centre for gynecological malignancies. METHODS: It was a retrospective descriptive study over a period of four years, from June 1(st. Koffi et al. presented that tumor grade II exemplified 58.4% of females, however grade I in addition II were in the comparable amount (20.8%) (22) which is a referral centre for gynecological malignancies. METHODS: It was a retrospective descriptive study over a period of

four years, from June 1(st. These results are conflicting to those reported by Meye et al. that, the grade II detained the past location after grade I and III (23) which is a referral centre for gynecological malignancies. METHODS: It was a retrospective descriptive study over a period of four years, from June 1(st.

According to association between ages groups and behavior of lesion either benign or malignant, there is significant association between age groups and cancer of breast, 37% of females in age group 41- 50 years old with malignant breast cancer, while 20% of malignancy in age group (31 – 40) and (51 – 60) years old. 13% of malignancy in age group 61 – 70 years old and only 6% in female over 71 years old age. Augustin et al. stated the same results that most patients in ages of 35 to 44 and 45 to 54 with 28.2% and 29.3% correspondingly. We can roughly that the major people found is dependable with some preceding studies approved globally (24, 25). Another study done by Hai-long Chen et al. showed that age groups from 50-59 years old was most group with breast cancer (26.4%), less than 30 years old and less than 40 years old represented low amount of patients (6.4%) (26). In another study stated that 6.4% of patients with breast cancer younger than 40 years old while 93.6% of them more than 40 years old, so the incidence of females to developed breast ca. was more in age group more than 40 years old so all women must do routinely breast cancer screening (27).

Conclusion

After pathological assessment of biopsies, malignant breast cancer (IDC/ grade II, IDC/ grade III, ILC and fibrocystic changes) most common changes, significant association between age group and breast cancer development more age group 41- 50 years old and then (31- 40), (51– 60) years old respectively.

Ethical Clearance: The Research Ethical Committee at scientific research by ethical approval of both MOH and MOHSER in Iraq

Conflict of Interest: Non

Funding: Self-funding

References

1. Shaukat U, Ismail M, Mehmood N. Epidemiology, major risk factors and genetic predisposition for breast cancer in the Pakistani population. *Asian Pac J Cancer Prev.* 2013;14(10):5625–5629.
2. Ly M, Antoine M, André F, Callard P, Bernaudin JF, Diallo DA. Breast cancer in Sub-Saharan African women. *Bull Cancer.* 2011;98(7):797–806.
3. Elgaili EM, Abuidriss D, Rahman M, Michalek AM, Mohammed SI. Breast cancer burden in central Sudan. *Int J Womens Health.* 2010;9(2):77–82.
4. Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, et al. LOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC Cancer Base No. 11 [Internet] Lyon, France. In: *International Agency for Research on Cancer.* globocan.iarc.fr; 2013.
5. Naderimagham S, Alipour S, Djalalinia S, Kasaeian A, Noori A, Rahimzadeh S, et al. National and sub-national burden of breast cancer in Iran; 1990–2013. *Arch Iran Med.* 2014;17(12):794–799.
6. Bray F, Ren JS, Masuyer E, Ferlay J. Global estimates of cancer prevalence for 27 sites in the adult population in 2008. *Int J Cancer.* 2013;132(5):1133–1145.
7. Majeed W, Aslam B, Javed I, Khaliq T, Muhammad F, Ali A, et al. Breast cancer: major risk factors and recent developments in treatment. *Asian Pac J Cancer Prev.* 2014;15(8):3353–3358.
8. Cole P, Mark Elwood J, Kaplan SD. Incidence rates and risk factors of benign breast neoplasms. *Am J Epidemiol.* 1978; 108:112–20.
9. Hutchinson WB, Thomas DB, Hamlin WB, et al. Risk of breast cancer in women with benign breast lesion. *J Natl Cancer Inst.* 1980; 65:13–20.
10. Mansel RE. Benign breast disease. *Practitioner.* 1982; 232:830–37.
11. Sainsbury RC. Bailey and Love's Short Practice of Surgery. 25th. London: Edward Arnold Ltd.; 2008. Breast In: Norman WS, Bulstrode CJK, P.Ronan O'Connel editors; pp. 827–35.
12. Sangma MB, Panda K, Dasiah S. A clinico-pathological study on benign breast diseases. *J Clin Diagn Res.* 2013;7(3):503–506.
13. Allred DC. Ductal carcinoma in situ: terminology, classification, and natural history. *J Natl Cancer Inst Monogr.* 2010; 2010: 134-138.
14. Erbas B, Provenzano E, Armes J, Gertig D. The natural history of ductal carcinoma in situ of the breast: a review. *Breast Cancer Res Treat.* 2006; 97: 135-144.
15. Sanders ME, Schuyler PA, Dupont WD, Page DL.

- The natural history of low-grade ductal carcinoma in situ of the breast in women treated by biopsy only revealed over 30 years of long-term follow-up. *Cancer*. 2005; 103: 2481-2484.
16. AMERICAN JOINT COMMITTEE ON CANCER. *AJCC Cancer Staging Manual - 8th Edition*. 2017. 1–1001 p.
 17. Balekouzou A, Yin P, Pamatika CM, et al. Epidemiology of breast cancer: retrospective study in the Central African Republic. *BMC Public Health*. 2016;16(1):1230. Published 2016 Dec 7.
 18. Yu, Z. G. *et al.* The prevalence and correlates of breast cancer among women in Eastern China. *PLoS ONE* 7, (2012).
 19. Spivey GH, Perry BW, Clark VA, Coulson AH, Coulson WF. Predicting the risk of cancer at the time of breast biopsy. Variation in the benign to malignant ratio. *Am Surg*. 1982 Jul;48(7):326–32.
 20. Ohene-Yeboah M, Adjei E. Breast cancer in Kumasi, Ghana. *Ghana Med J*. 2012 Mar;46(1):8–13.
 21. Essiben F, Foumane P, Mboudou ET, Dohbit JS, Mve Koh V, Ndom P. [Diagnosis and treatment of breast cancer in Cameroon: a series of 65 cases]. *Mali Med*. 2013;28(1):1–5.
 22. Koffi B, Sepou A, Doui DA, Goumba C, Djabanga S. Epidemiological and histological characteristics of breast cancers in Bangui [French article] *Med Afr Noire*. 2004;1(2):112–4.
 23. Meye JF, Ngomo KM, Diallo I. Breast cancer hospital in Libreville [Article in French] *Med Afr Noire*. 2004;51(8–9):479–72.
 24. Mobima T, Ouiamon R, Koandongui F, Glamende, Koffi B. Histo-sonographic correlation of breast lesions in Bangui (CAR) [French article] *J Afr Imag Med*. 2015;4(7):431–438.
 25. Adebamowo CA, Ayayi OO. Breast Cancer in Nigeria. *West Afr J Med*. 2000;19:179–181.
 26. Chen, H. L., Zhou, M. Q., Tian, W., Meng, K. X. & He, H. F. Effect of age on breast cancer patient prognoses: A population-based study using the SEER 18 database. *PLoS ONE* 11, (2016).
 27. Gnerlich JL, Deshpande AD, Jeffe DB, Sweet A, White N, Margenthaler JA. Elevated breast cancer mortality in women younger than age 40 years compared with older women is attributed to poorer survival in early-stage disease. *Journal of the American College of Surgeons*. 2009;208(3):341–7.