

Infection with Kala Azar Among Children in Babylon Province and Their Effect on Hematocrit Reading and Their Relation with Blood Type

Ahmed M. Al-Mosawy Lekaa A. AL Qurashi Zainab H. Al-Zubiady
Collage of dentistry/Babylon University

Abstract

A prospective study of Hospitalized cases of children in Babylon Maternity and Children Hospital , Al-Hilla General Hospital and Al-Hashmya Hospital was carried out from 1st September 2007 to 1st March 2008, blood samples were collected . It has been documented 54 patients of children were infected with *Leishmania donovani* (Kala azar) by positive serological test (IFAT) . The age of those children was between 3 months to 3 years old , they were divided into two groups (under 1 years old and 1-3 years old) . Hematocrit of these children were measured and compared with Hematocrit of other healthy children . Blood group test (ABO) was also carried out. Kala Azar was highly number in children within age 1-3 years old (46).Also Al-Hashmya district showed highly number of Leishmaniasis(32)than other districts including Al-Mahaweel , Al-Hilla and Ashomaly(9,7and6) respectively Leishmaniasis caused highly significant decrease ($p<0.01$) in hematocrit of children under one years old (27.13 ± 1.48) and highly significant decrease ($p<0.001$) in hematocrit of children within 1-3 years old (26.96 ± 0.39). Kala Azar was highly percentage in blood group type A (24.5%) as compared with other blood group type O,B and AB(24%,20.3%and12.9%)respectively.

Hematocrit		Blood group		IFAT
()	%	()	%	()
27.13	%	24.5	%	(P<0.01)
A	(,)	AB	(,)	(P<0.01)
B	(,)	O	(,)	(,)

Introduction

Visceral leishmaniasis or(kala azar) is a systemic disease which is caused by *Leishmania donovani* complex(Fakhar *et al.*, 2006) which is known to be endemic in areas of the middle east including Yemen , Oman , Kuwait , Iraq , Saudi Arabia , United Arab Emirates and Bahrain (Zeibig ,1997 ; Elnour *et al.*, 2001 ; Al-Marzoki , 2002 ; Al-Muhamadi *et al.*, 2004 ; Fakhar *et al.*, 2006) as well as in Tunis (Khaldi *et al.*, 1991) and Turkey (Ozensoy *et al.* 1998) . In Europe visceral leishmaniasis are endemic in all Mediterranean countries such as southern France (Mindoir *et al.*, 1998) Southern Greece (Maltezou *et al.*, 2000) and Southern Spain (Pineda *et al.*, 1998) Also there is documented cases of kala azar in central European countries such as Germany (Bogdan *et al.*, 2001)

Materials and Methods

Hospitalized cases of children in Babylon Maternity and Children Hospital , Al-Hilla General Hospital and Al-Hashmya Hospital was carried out from 1st

September 2007 to 1st March 2008 . It has been diagnosed 54 patients of children were infected with *Leishmania donovani* by demonstrate *Leishmania* antibody in their sera by using indirect Immunofluorescent Antibody Technique (IFAT) according to Randox Co. kit ,as recommended by manufacture.

The age of those children was between 3 months to 3 years old , which arranged into two groups . First group from 3 months to under one year . Second group from one year to three years old (Al-Muhammadi *et al.*, 2004) .

Blood was collected by antecubital veinpuncture at 9–11 a.m. Hematocrit (Hct) of these children was measured according to Talib & Khurana (1996) , and compared with Hct of other healthy children (control 5 children of each age group) . Blood group ABO test was carried out according to Talib & Khurana (1996). Results of Hct were statistical analyzed using t-test (Campbell , 1967).

Results and Discussion

Table (1) shows highly number of visceral leishmaniasis in children within 1-3 years old(46) as compared with the number of infection in children under one years old (8) which agree with the results of Mindoir *et al.*(1998) . This goes with the fact that the infantile type is the commonest type in our area (Al-Marzoki , 2002) and may be the symptoms can be persist for weeks to several months before patients come to medical attention (Ravanbod, 2002).

There was no such difference in the sex factor (male: female 31:23) of visceral leishmaniasis in children which agree with Al-Marzoki (2002) and Al-Muhammadi *et al.*(2004) .

In table (2) , Al-Hashmya district showed highly number of kala azar infection (32) than other districts of Babylon province including AL-Mahaweel, Al-Hilla and Ashomaly (9,7and6) respectively, this agree with Al-Marzoki (2002) and Al-Muhammadi *et al.*(2004) .

Iraq has been reported to be one of the endemic area of Kala azar (Zeibig ,1997; Al-Marzoki , 2002 ; Al-Muhamadi *et al.*, 2004). The high prevalence of leishmaniasis in Al-Hashmya district and then Al-Mahaweel district may be due to the highly prevalence of the vector (Sandflies) of visceral *Leishmaniasis* in these specific area that get a big attention for controlling in this area .

Visceral leishmaniasis caused highly significant decrease ($p<0.01$) in Hct of children under one years old (27.13 ± 1.48) and very high significant decrease ($p<0.001$) in Hct of children within 1-3years old (26.96 ± 0.39) as shown in table (3) . This result agree with Al-Muhammadi *et al.*(2004) . That is to say visceral leishmaniasis caused anemia (Mindoir *et al.*, 1998; Ozensoy *et al.*, 1998; Maltezou *et al.*, 2000; Elnour *et al.*, 2001 ; Al-Muhammadi *et al.*, 2004 ; Joshi *et al.*, 2006) . Anemia always present with leishmaniasis and may be severe , it is usually normocytic normochromic and appears due to a combination of factors including hemolysis , marrow replacement with visceral *Leishmaniasis* infected macrophages , hemorrhage , splenic sequestration of erythrocytes , hemodilution (Ravanbod,2002) and due to effect of malnutrition (Al-Marzoki , 2002) then it appears as iron deficiency (Microcytic hypochromic) anemia (Elnour *et al.*, 2001). In table (4)visceral leishmaniasis was highly percentage in blood group type A (42.5%) as compared with other blood group types O, B, and AB (24%,20.3%and12.9%) respectively This was agree with (Al-Mamouri , 2000) who found that protozoa related with type A blood group due to the presence of genetic factors and the antigens of these protozoa are similar to the antigens of type A blood group subjects and so do not recognized by immune system.

Table (1) : Distribution of Leishmaniasis according to age and sex .

Age(year)	Male	percentage	Female	percentage
Less than 1	3	5.5	5	9.2
1-3	28	15.8	18	33.3
Total	31	57.4	23	42.5

Table (2) : Distribution of leishmaniasis according to location .

No	Location		Patient	percentage
1	Al-Hilla District	Al-Hilla center Al-Tuhmazya	7	12.9%
2	Al-Mahaweel District	Al-Mahaweel center Jibalah Al-Neel center	9	16.6%
3	Al-Hashmya District	Al-Hashmya center Al-Qassim Al-Midhatya	32	59.2%
4	Ashomaly District		6	11.1%
	Total		54	100

Table (3) : Effect of leishmaniasis on hematocrit reading .

Age(year)	Hematocrite reading (percentage)	
	Control	Patient
Less than 1	34 ± 0.39	27.13 ± 1.48*
1 – 3	36.2 ± 1.48	26.96 ± 0.39**

* Significant decrease (p<0.01)

** Significant decrease (p<0.001)

Table (4) : Relation of leishmaniasis with blood group (ABO).

Blood group	No.of patients	percentage
A	23	42.5
B	11	20.3
O	13	24.0
AB	7	12.9
Total	54	100

Recommendation

From above we recommend

1. The screening (Survey) study should be taken on the areas close to those documented endemic areas in this study.
2. Protection against sandflies accomplished by repellents, protective clothing are essential measures to reduce future *L. donovani* infection. Promote treatment of human infections as well as control of sandflies population will also help to control of disease .

References

- Al-Mamouri, A. K. (2000). Epidemiology of intestinal parasites and head lice in pupils of some primary schools at Al-Mahaweel district, Babylon province. MSc. Thesis , Sci. Coll. , Babylon Univ. : 122pp. (In Arabic).
- Al-Marzoki, J.M. (2002) . Clinical and laboratory study of kala azar in Hilla , Iraq. Med. J. Babylon 7(4):1715-1721.
- Al-Muhammadi , M.O. Shujini, G.S. and Noor ,M.H. (2004). Hematological changes in children suffering from visceral Leishmaniasis (kala azar) . Medical Journal of Babylon ,1 (3,4) :322-29.
- Bogdan, C.; Schönian, G.; Bañules, A. , Hide, M. ; Pratlong, F. ; Lorenz, E. ; Röllinghoff, M. & Mertens, R. (2001). Visceral Leishmaniasis in a German Child Who Had Never Entered a Known Endemic Area: Case Report and Review of the Literature. CID.32:3002-6.
- Campbell, R.C. (1967). Statistics for biologists .Cambridge Univ. Press: 242 pp.
- Elnour, I.B. ; Akinbami , F.O. ;Shaked, A. and Venugopalan ,P.(2001) Visceral Leishmaniasis in omani children : a review Ann. Trop. Pediatr. Jun; 21 (2): 159-63
- Fakhar, M; Asgari, Q; Motazedin, M.H. ; Moheballi, M; Hatam, Gh.R. and Alborzi, A.V. (2006). First report of kala azar from Qeshm Iland in Persian Gulf . Iranian J Parasitol, 1(1): 53-56.
- Joshi, S.; Bajracharya, B. L. & Baral, M. R. (2006) . Kala-azar (Visceral Leishmaniasis) from Khotang . Kathmandu Univ. Med. J. 4(2): 232-234
- Kaldi ,F. ;Achouri ,E.;Gharbi ,A.; Debbabi ,A. and Ben Naceur ,B.(1991) Visceral Leishmaniasis in children. Astudy of hospitalized cases from 1974 to 1988 at children's hospital in Tunis. Med. Trop. 51(2) : 143-8. (Abst. Article in French).
- Maltizou, H.C.; Saifas, M.; Mavrikou ,M.; Spyridis, P.; Stovrinadis, C.; Karpathios , Th. And kafetzis, D.A. (2000). Visceral Leishmaniasis during childhood in southern Greece. 31:1139-1143
- Mindoier, P.; Piarrous, R.; Garnier, J.M. ,Unal, D.;perrimond, H. and dumon ,H. (1998). Pediatric Visceral Leishmaniasis in Southern France. Pediatr .Infect. Dis. J. 17(8) : 701-4. (Abst.)
- Ozensoy, S.; Ozbel, Y. ; Turjay, N.; Alkan, M.Z.; Gul, K; Gilman-Sachs, A; Chang, K.P. ;Reed, S.G. and Ozcel ,M.A.(1998). Serodiagnosis and epidemiology of visceral Leishmaniasis in Turkey. Am.J. Trop. Hyg. 59(31:363-9.(Abst)
- Pineda, J.A.;Gallardo, J.A.; Macias ,J; Delgado, J.;Regordan, J.; Regordá, C.; Morillas , F; Relimpio, F; Martin-Sánchez,J.; Sánchez-Quijano, A, ; Leal,M.& Lissen, E. (1998) . Prevalence of and factors associated with visceral leishmaniasis in human immunodeficiency virus type 1- infected patients in

Southren Spain J. Clin. Microbio. 36(9):2419-2422.

Ravanbod, M. (2002). Kala Azar in adults ; a case presentation and review. SEMJ 3(4).

Talib, V.H. & Khurana, S.R. (1996). A handbook of medical laboratory technology. 5th, edn., C.B.S. Publ., New Delhi.

Zeibig, E. A. (1997). Clinical parasitology: A practical approach. W. B. Saunders Co., Philadelphia: 325 pp.