

JOURNAL OF THE UNIVERSITY OF KARBALA

An Academic Journal Adopted for Scientific Promotions Issued by
Karbala University



Editor in chief

Prof. Dr. Hassan Aoda Al-Ghanimi

Secretary of Editor

Assis.Prof.Dr. Shafi Hussein Al-Sharifi

Director Editor

Muneer Abdul-Ali Al-Da'amie

Technical Office

Assel A. Hussien

Enaam G. Radi

Journal of Kerbala University , Vol.4. No.1.Scientific . 2008

18.	<p align="center">Design and Implementation of an Access Control System for a Bank</p> <p align="center">المدرس المساعد/عباس فاضل محمد علي قسم علوم الحاسبات- كلية العلوم - جامعة كربلاء المبرمج/بشرى جابر محمد جواد قسم ادارة الاعمال - كلية الادارة والاقتصاد - جامعة كربلاء</p>	159-169
19.	<p align="center">Kriging of Groundwater Level - A Case Study of Dibdiba Aquifer in Area of Karbala-Najaf</p> <p align="center">Assist. Teacher –Waqed Hameed Al-Mussawi Department of Computers - College of Science- Karbala University</p>	170-182
20.	<p align="center">Study of some topological concepts in bitopological spaces</p> <p align="center">يزي كاظم مندي الطالقاني سكينة عبد الله ليلو جامعة بابل كلية التربية/قسم الرياضيات</p>	183-188
21	<p align="center">Some Hematological and Histological changes in seminiferous tubules of pigeon (<i>Columba livia</i>) that exposed to different periods of light</p> <p align="center">Sinaa Al_Bazii Kerbala university, Education college Biological department</p>	189-195
22.	<p align="center">Active Inhibitor as Corrosion Protective of Carbon Steel</p> <p align="center">Dr. Ali Hubi Haleem Collage of Materials Engineering-Babylon University</p>	196-207
23.	<p align="center">Effect of Mineral Filler and Additives Materials on the Adhesion Properties between Asphalt and Aggregate</p> <p align="center">Dr. Abdul Haq H. Alhaddad Highway and Transportation Department – College of Engineering – Al- Mustansiriya University</p>	208-217
24.	<p align="center">Effect of Rizolex on <i>Rhizoctonia solani</i> Kuhn isolates and biocontrol fungus <i>Trichoderma harzianum</i> Rifai <i>in vitro</i></p> <p align="center">FAKHIR R. HAMEED COLLEGE OF AGRICULTURE –UNIVERSITY OF BABYLON</p>	218-223
25.	<p align="center">THE MAXIMUM NUMBER IN WHICH EVERY STRONG TOURNAMENT CONTAINS A TRANSITIVE SUBTOURNAMENTS</p> <p align="center">Dr . Nihad ABDEL – JALIL MAJID / Assistant professor Dr_ nihad45@yahoo.com. Dep of Mathematics</p>	224-227
26.	<p align="center">The General Solution Of the Partial Differential Equation which has The General form</p> <p align="center">$P(x,y)Z_x + Q(x,y)Z_y + R(x,y,z)Z = G(x,y,z)Z^n, n \neq 1$</p> <p align="center">Assis.prof.Ali Hassan Mohammed Kufa University. College of Education. Department of Mathematics Assis. Lect. Rehab Ali Kuder Kufa University. College of Education. Department of Computer Sciences</p>	228-232

Study of some topological concepts in bitopological spaces

دراسة بعض المفاهيم التبولوجية في الفضاء ثنائي التبولوجي
 يزي كاظم مهدي الطالقاني
 سكيمة عبد الله ليلو
 جامعة بابل /كلية التربية/قسم الرياضيات



Abstract:

A new definition of bitopological space is introduced in this paper with its separation axiom and continuity. Connected and compact set are defined with some theorems and also $T_{1,2}$, semi $T_{1/2}$, semi $T_{1/3}$, spaces are defined in this paper.

ملخص البحث :

يقدم هذا البحث تعريف جديد للمجموعة المفتوحة في الفضاء ثنائي التبولوجي (X, P_1, P_2) في هذا الفضاء عرفنا الترابط والترانس والإستمرارية وقمنا بدراسة بعض العلاقات بينها، بالإضافة إلى ذلك قمنا بتعريف فضاء $T_{1/2}$ وفضاء $semiT_{1/2}$ وفضاء $semiT_{1/3}$ في فضاء ثنائي التبولوجي ودراسة بعض خواصها فيه.

1- introduction :

A bitopological space (X, p_1, p_2) [J.C.Kelly "bitopological spaces", 1963] is a non-empty set X with two topologies p_1 and p_2 on X .

[Caldas, M, "semi- $T_{1/2}$ -space" 1994] defined the concept of a new class of topological spaces called semi- $T_{1/2}$ spaces.

[Kumar, "Between semi closed set and semi- ρ -closed set" 1991] introduced a new class of spaces namely semi- $T_{1/3}$ spaces he proved that the class of semi $T_{1/3}$ spaces properly contains the class of semi- $T_{1/2}$ spaces and he defined a new map namely ψ -continuity and he discussed the relation between this map and semi-continuity [Biswas, N "characterization of semi-continuous function", 1970], g -continuity [Caldas, M, "on g -closed sets and g -continuous mapping", 1993], [R.Devi, H.Maki and K.Balachandran, "semi-generalized closed map and generalized closed map", 1993], [K.Balachandran, P.Sundaram and H.Maki, "semi-generalized continuous maps and semi- $T_{1/2}$ ", 1991], sg -continuity [P.sundaram, H.Maki, K. Balachandran, "on generalized continuous maps in topological spaces", 1991], [P.Bhattacharya and B.K.Lahiri, "semi-generalized closed sets in topology" 1987] and gs -continuity [Miguel Caladas Cueva and Ratnesh Kumar saraf, "A research on characterization of semi- $T_{1,2}$ -spaces" 2000].

The purpose of this paper is to give a new definition of these concepts in Bitopological spaces.

2- Basic definitions and theorems

We would like to point out that all the definitions provided in this research have been formulated by researchers by adoption of their counterparts in topological spaces.

Definition (2-1)

Let (X, p_1, p_2) be a bitopological space then a subset A of X is said to be open iff there exists T_i - open set U such that $U \subseteq A$ and $\bigcap cl_{p_i}(U) \subseteq A, i=1,2$, this open set denoted by δ -open set

Example(2-2):

let $X = \{a, b, c, d\}$, $T_1 = \{X, \phi, \{a\}, \{b\}, \{a, b\}\}$, $T_2 = \{X, \phi, \{a\}, \{c\}, \{a, c\}\}$ then δ -open set = $\{X, \phi, \{a, b, d\}, \{b, c, d\}, \{a, c, d\}\}$

Remark(2-3)

The intersection of two δ -open sets is not necessarily δ -open while the union is δ -open set

Proof: let $\{A_\lambda : \lambda \in \Lambda\}$ be any arbitrary collection of δ -open set, then there exist T_i -open set U_λ such that $U_\lambda \subseteq A_\lambda$ and $\bigcap cl_{p_i}(U_\lambda) \subseteq A_\lambda, i=1,2$ for each λ .

Since

$$\square_{\lambda \in \Lambda} (\bigcap_{i=1,2} cl_{p_i}(U_\lambda))$$

$$= (\text{cl}_{p_1}(\square(U_\lambda)) \cap (\text{cl}_{p_2}(\square(U_\lambda))) = \bigcap_{i=1,2} \text{cl}_{p_i}(\square_{\lambda \in \Lambda}(U_\lambda)) \subseteq \square_{\lambda \in \Lambda} A_\lambda$$

Remark (2-4)

- The set of all ∂ -open sets is not a topological space.
- if a is p_i -closed set for $i=1,2$ then A is ∂ -open set

Example(2-5):

Let $X=\{a,b,c,d\}$ and $T=\{X, \phi, \{a\}, \{b,c\}\}$, $T=\{X, \phi, \{a\}, \{c,d\}\}$

∂ -open set $=\{X, \phi, \{a\}, \{a,b\}, \{a,c\}, \{a,d\}, \{a,b,c\}, \{a,b,d\}, \{b,c,d\}\}$

Since $\{a,b,d\} \cap \{c,b,d\} = \{b,d\}$ which is not ∂ -open set then the set of all ∂ -open sets is not topological space .

Notation (2-6)

let (X, p_1, p_2) be a bitopological space and let Y be a subset of X then a subset B of Y is said to be ∂_Y -open (the open set in (Y, p_{1Y}, p_{2Y})) iff there exist p_{iY} open set W such that $W \subseteq B$ and $\bigcap \text{cl}_{p_{iY}}(W) \subseteq B$

Theorem (2-7)

if A is ∂ -open set and B is ∂_Y -open set where Y is a subset of X which is not ∂ -open set then $A \cap B$ is ∂_Y -open set and $A \cup B$ is ∂ -open set

proof:- since A is ∂ -open set then there exist T_i -open set U such that $U \subseteq A$ and $\bigcap \text{cl}_{p_i}(U) \subseteq A$ and since B is ∂_Y -open set then there exist T_{iY} -open set W such that $W \subseteq B$ and $\bigcap \text{cl}_{p_{iY}}(W) \subseteq B$. then $[\bigcap \text{cl}_{p_i}(U)] \cap [\bigcap \text{cl}_{p_{iY}}(W)] \subseteq A \cap B$ and then $\text{cl}_{p_1}(U \cap W) \cap Y \cap \text{cl}_{p_2}(U \cap W) \cap Y \subseteq (\text{cl}_{p_1}(U) \cap \text{cl}_{p_1}(W)) \cap Y \cap (\text{cl}_{p_2}(U) \cap \text{cl}_{p_2}(W)) \cap Y \subseteq A \cap B$, from the last statement we get the result .

$$\begin{aligned} & [\bigcap \text{cl}_{p_i}(U)] \cap [\bigcap \text{cl}_{p_{iY}}(W)] \subseteq A \cap B, \text{ then} \\ & (\text{cl}_{p_1}(U) \cap \text{cl}_{p_2}(U)) \cap (\text{cl}_{p_{1Y}}(W) \cap \text{cl}_{p_{2Y}}(W)) \\ & = (\text{cl}_{p_1}(U) \cap \text{cl}_{p_{1Y}}(W)) \cap (\text{cl}_{p_1}(U) \cap \text{cl}_{p_{2Y}}(W)) \cap (\text{cl}_{p_2}(U) \cap \text{cl}_{p_{1Y}}(W)) \cap (\text{cl}_{p_2}(U) \cap \text{cl}_{p_{2Y}}(W)) \\ & = [\text{cl}_{p_1}(U) \cap W] \cap [\text{cl}_{p_2}(U) \cap W] \cap [\text{cl}_{p_1}(U) \cap Y] \cap [\text{cl}_{p_2}(U) \cap Y] \\ & \cap [\text{cl}_{p_1}(U) \cap \text{cl}_{p_{2Y}}(W)] \cap [\text{cl}_{p_2}(U) \cap \text{cl}_{p_{1Y}}(W)] \subseteq A \cap B \text{ and then} \\ & \bigcap \text{cl}_{p_i}(U \cap W) \subseteq A \cap B . \end{aligned}$$

Remark (2-8)

By theorem (1-5) we can define the subspace Y of a topological space X as follow:
if Y is a subset of X such that Y is ∂ -open set then $\partial_Y\text{-open} = \{G \cap Y : G \text{ is } \partial\text{-open set in } X\}$

Notation (2-9)

Let $(X, p_1, p_2), (Y, w_1, w_2)$ are two bitopological spaces , a function $f: (X, p_1, p_2) \rightarrow (Y, w_1, w_2)$ is said to be ∂ -continuous iff $f^{-1}(V)$ is ∂ -open(∂ -closed) set in X for each V is ∂ -open(∂ -closed) set in Y

Theorem (2-10)

- A function $f: (X, p_1, p_2) \rightarrow (Y, w_1, w_2)$ is said to be
- 1- ∂ -open mapping iff $f(U)$ is ∂ -open whenever U is ∂ -open set in X
 - 2- ∂ -closed iff $f(U)$ is ∂ -closed in Y whenever U is ∂ -closed set in X
 - 3- ∂ -bi continuous iff f is ∂ -open and ∂ -continues
 - 4- ∂ -homeomorphism iff f is bijection mapping , ∂ -continuous and f^{-1} is ∂ -continuous

Proof: we using the same proof in [1] with replacing the open set(closed set) by ∂ -open(∂ -closed) set respectively.

Notation (2-11)

Let (X, p_1, p_2) be a bitopological and A,C are two subsets of X we say that A and C are separated iff $A \cap \text{cl}_{p_i}(C) = \emptyset$ and $C \cap \text{cl}_{p_i}(A) = \emptyset$ for $I=1,2$

Remark(2-12)

If A ,C are p_i -separated, for $I=1$ or 2 then it is not necessary that A,C are separated in (X, p_1, p_2)

Example(2-13)

In example(2-5) $\{a\}$ and $\{b,c\}$ which are T_1 -separated but are not separated in (X,p_1,p_2)

Theorem(2-14)

let Y be a subspace of a bitopological space X such that Y is ∂ -open set and let A,C be two subset of Y . then A, C are π -separated if and only if they are π_Y -separated.

Proof: by using the relation $\text{clip}(A) \cap Y = \text{clip}_Y(A)$ and $\text{clip}(C) \cap Y = \text{clip}_Y(C)$ the result exist

Notation(2-15)

Let (X,p_1,p_2) be a bitopological space . a subset A of X is said to be ∂ -disconnected if and only if it is the union of two non-empty π -separated subsets for $I=1,2$. A is said to be ∂ -connected iff it is not ∂ -disconnected .

Example(2-16)

$X=\{a,b,c\}$ $T_1=\{X,\phi,\{a\},\{b,c\}\}$ $T_2=\{X,\phi,\{a,c\}\}$ then X is ∂ -disconnected and $\{a,c\}$ is ∂ -connected

Theorem(2-17)

Let Y be a subspace of a bitopological space X and A subset of Y such that Y is ∂ -open set then A is ∂ -disconnected iff it is ∂_Y -disconnected and A is said ∂ -connected iff it is ∂_Y -connected

Proof: by theorem (2-14) A is union of π -separated sets iff it is π_Y -separated sets and hence the result .

Notation (2-18)

Let (X,p_1,p_2) be a bitopological space and A be a subset of X is said to be compact iff every π -open cover of A has π -finite sub cover for $I=1$ or 2

Example(2-19)

Let $X=\{a,b,c\}$, $T_1=\{X,\phi,\{a\},\{c\},\{a,c\}\}$, $T_2=\{X,\phi,\{a\},\{b,c\}\}$ then (X,p_1,p_2)

Is compact with respect to T_1 or T_2 .

Remark(2-20)

If (X,p_1) or (X,p_2) is compact then (X,p_1,p_2) is compact

Proof : by the definition of compactness in this paper clearly that if we take any open cover of X then this cover we take from p_1 or p_2 and hence the open cover containing a finite sub cover.

Theorem(2-21)

Let Y be subspace of a bitopological space X such that Y is ∂ -open set and let $D \subset Y$. then D is compact relative to X if and only if D is compact relative to Y .

Let D is compact relative to X and $\{G_\lambda : \lambda \in \Lambda\}$ is π_Y -open cover of A in Y then $A \subseteq \bigcup \{G_\lambda : \lambda \in \Lambda\}$, then there exist V_λ which is π -open sets in X such that $G_\lambda = V_\lambda \cap Y$ for every $\lambda \in \Lambda$ and then $D \subseteq \bigcup \{V_\lambda : \lambda \in \Lambda\}$ then $\{V_\lambda : \lambda \in \Lambda\}$ is open cover of D in X and since D is compact relative to X then there exist $\lambda_1, \lambda_2, \dots, \lambda_n$ such that

$D \subseteq V_{\lambda_1} \cup V_{\lambda_2} \cup \dots \cup V_{\lambda_n}$ and then $D \subseteq [V_{\lambda_1} \cup V_{\lambda_2} \cup \dots \cup V_{\lambda_n}] \cap Y = G_{\lambda_1}$ and there for D is compact relative to Y . and in the same way we proof the converse.

Theorem(2-22)

Let (X,p_1,p_2) be a compact bitopological space then a π -closed F of X is compact for $i=1$ or 2 respectively. subset

Proof:

Let $C=\{G_\lambda : \lambda \in \Lambda\}$ is π -open cover of F where F is π -closed set in X let

$D=\{G_\lambda : \lambda \in \Lambda\} \cup \{X-F\}$ form π -open cover of X since X is compact the D has a finitely sub collection of D covers X and then covers F , and then F is compact.

Notation (2-23)

a bitopological space (X,p_1,p_2) is say to be

1- ∂ - T_0 iff for each x, y in X thee exist ∂ -open set U such that $x \in U, y \notin U$

2- ∂ - T_1 iff for each two distinct point x,y there exist two ∂ -open sets U,W such that $x \in U, y \in W$

- 3- ∂ - T_2 space iff for each two distinct point x, y there exist two ∂ -open sets U, W such that $x \in U, y \notin W$ and $U \cap W = \emptyset$
- 4- ∂ - $T_{2 1/2}$ space iff any two points of X can be separated by ∂ -closed neighborhood
- 5- ∂ -regular iff for each ∂ -closed set F and $y \notin F$ there exist two ∂ -open sets U, W such that: $F \subseteq U, y \in W$ and $U \cap W = \emptyset$
- 6- ∂ - T_3 iff it is ∂ - T_1 and regular
- 7- ∂ -completely regular space if and only if for an ∂ -closed set F in X and x any point in X not in F there exist ∂ -continuous map $f: X \rightarrow [0, 1]$ such that $f\{F\} = \{1\}, f(X) = 0$
- 8- ∂ - $T_{3 1/2}$ space iff this space is ∂ -completely regular and ∂ -hausdorff
- 9- ∂ -normal iff for each two disjoint π -closed sets F, H for $i=1, 2$ there exist two ∂ -open set U, W such that $F \subseteq U, H \subseteq W$ and $U \cap W = \emptyset$
- 10- T_4 space if and only if X is ∂ - T_1 and ∂ -normal space
- 11- completely normal space if and only if for a separated sets A, C there exist two ∂ -open sets U, W such that $A \subseteq U$ and $C \subseteq W, U \cap W = \emptyset$ and we denoted this space by $[CN]$
- 12- ∂ - T_5 space iff it is both ∂ -completely normal and hausdorff

Theorem(2-24)

Let (X, p_1, p_2) be a bitopological space then every ∂ -complete normal space is ∂ -normal

Proof:

Let A, B are two disjoint π -closed sets then $A \cap \text{cl}_{\pi}(B) = \emptyset$ and $B \cap \text{cl}_{\pi}(A) = \emptyset$ and there for A, B are two disjoint separated sets in X and since X is ∂ -complete normal space there exist two disjoint ∂ -open set U, W such that $A \subseteq U$ and $B \subseteq W$, then (X, p_1, p_2) is ∂ -normal space.

Theorem (2-25) let (X, p_1, p_2) be a bitopological space and Y be a subset of X such that $Y \in \partial$ -open(X) then ∂ - T_0, ∂ - T_1, ∂ - T_2, ∂ -regular, ∂ - T_3, ∂ - T_5 are hereditary property

Proof: see [1] with replacing every open set (closed set) by ∂ -open set (∂ -closed set) respectively .

3- new main results

Notation (3-1)

Let (X, p_1, p_2) be a bitopological space then a subset A of X is said to be

a- semi-open set if $A \subseteq \text{cl}_{\pi}(\text{int}_{\pi}(A))$ and semi-closed if $\text{int}_{\pi}(\text{cl}_{\pi}(A)) \subseteq A$ for $i=1$ or 2 .

b- generalized closed set (briefly g-closed set) iff $\text{cl}_{\pi}(A) \subseteq U$ where $A \subseteq U$ and U is ∂ -open set in (X, p_i) for $i=1$ or 2 .

c- semi-generalized closed (briefly sg-closed set) if $\text{scl}_{\pi}(A) \subseteq U$ whenever $A \subseteq U$ and U is semi-open set in (X, p_i) for $i=1$ or 2

d- generalized semi-closed (briefly gs-closed) iff $\text{scl}_{\pi}(A) \subseteq U$, where $A \subseteq U$ and U is ∂ -open set in (X, p_i) for $i=1$ or 2 .

e- ψ -closed set if $\text{scl}_{\pi}(A) \subseteq U$ whenever $A \subseteq U$ and U is sg-open set in (X, p_i) for $i=1$ or 2

Example(3-2)

Let $X = \{a, b, c\}$ and $T_1 = \{X, \emptyset, \{a, b\}\}$ $T_2 = \{X, \emptyset, \{a\}, \{b, c\}\}$ then $\{a, c\}$ is ψ -closed

And $\{b\}$ is sg-open set and sg-closed set but $\{b, c\}$ is not ψ -closed set .

proposition (3-3)

A subset A of X is semi-open set (respectively, semi-closed, sg-closed, gs-closed, g-closed, ψ -closed) in (X, p_i) for $i=1$ or 2 then it is semi-open set (respectively, semi-closed, sg-closed, gs-closed, g-closed, ψ -closed) in (X, p_1, p_2)

proof:

let A is π -semi-open (respectively, π -semi-closed, π -sg-closed, π -gs-closed, π -g-closed, π - ψ -closed) set then $A \subseteq \text{cl}_{\pi}(\text{int}_{\pi}(A))$ [respectively, $\text{int}_{\pi}(\text{cl}_{\pi}(A)) \subseteq A, \text{scl}_{\pi}(A) \subseteq U$ where $A \subseteq U$ and U is π - ∂ -open set, $\text{scl}_{\pi}(A) \subseteq U$ where $A \subseteq U$ and U is π -semi-open set, $\text{cl}_{\pi}(A) \subseteq U$ where $A \subseteq U$ and U is π - ∂ -open set, $\text{scl}_{\pi}(A) \subseteq U$ whenever $A \subseteq U$ and U is π -sg-open set] and then A is semi-open set (respectively, semi-closed, sg-closed, gs-closed, g-closed, ψ -closed) in (X, p_1, p_2) .

Notation (3-4)

A bitopological space (X, p_1, p_2) is said to be

a- $T_{1/2}$ space if every g -closed set is ∂ -closed set

b-Semi- $T_{1/2}$ space [Caldas, M., 1994] if every sg -closed set is semi-closed set

c- semi- $T_{1/3}$ space [M.K.R.S.Veera Kumar, 1991] if every ψ -closed set in it is semi-closed

Example(3-5)

Let $X = \{a, b, c\}$ and $T_1 = \{X, \phi, \{a\}, \{b, c\}\}$ $T_2 = \{X, \phi, \{b\}, \{b, c\}\}$ then (X, p_1, p_2) is semi- $T_{1/3}$ space and semi $T_{1/2}$ space

proposition (3-6)

If (X, p_i) is semi- $T_{1/3}$ space (respectively $T_{1/2}$ space, semi- $T_{1/2}$ space) for $i=1$ or 2 then (X, p_1, p_2) is semi- $T_{1/3}$ space (respectively $T_{1/2}$ space, semi- $T_{1/2}$ space)

proof: let A be a ψ -closed set (respectively, g -closed set, sg -closed set) in (X, p_1, p_2) by theorem (3-5) A is ψ -closed set (respectively, g -closed set, sg -closed set) in (X, p_i) , $i=1$ or 2 and since (X, p_i) is semi $T_{1/3}$ space (respectively $T_{1/2}$ space, semi- $T_{1/2}$ space) A is semi-closed set (respectively, ∂ -closed set, semi closed set) then (X, p_1, p_2) is semi $T_{1/3}$ space (respectively, $T_{1/2}$ space, semi $T_{1/2}$ space).

Remark:(3-7)[7]

1-every semi closed set and thus every ∂ -closed set is ψ -closed set

2- every ψ -closed [M.K.R.S.Veera Kumar, 1991] is sg -closed set and also gs -closed set 3- every semi-closed set is sg -closed set

proposition (3-8)

every semi- $T_{1/2}$ space is semi- $T_{1/3}$ space

proof:- let A be a ψ -closed set in X , by remark (3-7) A is sg -closed set and since X is semi $T_{1/2}$ space A is semi closed set and then (X, p_1, p_2) is semi $T_{1/3}$ space.

the converse is not true as we show in the following example

Example(3-9)

let $X = \{a, b, c\}$, $p_1 = \{X, \emptyset, \{a\}, \{b, c\}\}$, $p_2 = \{X, \emptyset, \{a, b\}\}$

Then (X, p_1, p_2) is not semi- $T_{1/2}$ space since $\{a, c\}$ is sg -closed but not semi closed set, however (X, p_1, p_2) is semi $T_{1/3}$ space.

Definition (3-10)[6]

For any subset E of (X, p_1, p_2) , $scl_{p_i}^*(E) = \bigcap \{A : E \subseteq A \text{ such that } A \in sd(X, p_1, p_2)\}$ where $sd(X, p_1, p_2) = \{A : A \subseteq X \text{ and } A \text{ is } sg\text{-closed in } (X, p_1, p_2)\}$ and $SO(X, p_1, p_2)^* = \{B : scl_{p_i}^*(B^c) = B^c\}$ for $i=1$ or 2 .

Proposition (3-11)

A bitopological space (X, p_1, p_2) is a semi- $T_{1/2}$ -space if and only if

$SO(X, p_1, p_2) = SO(X, p_1, p_2)^*$.

Proof

Since the semi closed sets and the sg -closed sets are coincide by assumption $scl_{p_i}(E) = scl_{p_i}^*(E)$ holds for every subset E of (X, p_1, p_2) there for we have $SO(X, p_1, p_2) = SO(X, p_1, p_2)^*$

Conversely let A is sg -closed set of (X, p_1, p_2) . then we have $A = scl_{p_i}^*(A)$ and hence $A^c \in SO(X, p_1, p_2)$ thus A is semi closed set there for (X, p_1, p_2) is semi $T_{1/2}$ space

Proposition (3-12)

A bitopological space (X, p_1, p_2) is semi $T_{1/2}$ space if and only if for each $x \in X$, $\{x\}$ is semi open or semi closed

Proof

Suppose that for some $x \in X$, $\{x\}$ is not semi closed. since X is the only semi open set containing $\{x\}^c$, the set $\{x\}^c$ is sg -closed set so it is semi closed set in the semi $T_{1/2}$ space (X, p_1, p_2) , therefore $\{x\}$ is semi open set

Conversely, since $SO(X, p_1, p_2) \subseteq SO(X, p_1, p_2)^*$ holds by theorem (3-11) it is enough to prove that $SO(X, p_1, p_2)^* \subseteq SO(X, p_1, p_2)$. let $E \in SO(X, p_1, p_2)^*$. Suppose that $E \notin SO(X, p_1, p_2)$. then $scl_{p_i}^*(E^c) = E^c$ and $scl_{p_i}(E^c) \neq E^c$ hold. There exist a point x of X such that $x \in scl_{p_i}(E^c)$ and $x \notin E^c (= scl_{p_i}^*(E^c))$. since

$x \notin \text{scl}_{p_i}(E^c)$ there exist sg-closed set A such that $x \notin A$ and $E^c \subset A$. by the hypothesis the singleton $\{x\}$ is semi-open set or semi-closed set.

Now if $\{x\}$ is semi-open set, since $\{x\}^c$ is semi closed set with $E^c \subset \{x\}^c$, we have $\text{scl}_{p_i}(E^c) \subset \{x\}^c$, i.e, $x \notin \text{scl}_{p_i}(E^c)$. this contradicts the fact that $x \in \text{scl}_{p_i}(E^c)$. therefore $E \in \text{SO}(X, p_1, p_2)$.

If $\{x\}$ is semi closed set, since $\{x\}^c$ is semi-open set containing the sg-closed set A ($\supset E^c$) we have $\text{scl}_{p_i}(E^c) \subset \text{scl}_{p_i}(A) \subset \{x\}^c$. therefore $x \notin \text{scl}_{p_i}(E^c)$. this is contradiction. therefore $E \in \text{SO}(X, p_1, p_2)$.

Hence in both cases we have $E \in \text{SO}(X, p_1, p_2)$, i.e, $\text{SO}(X, p_1, p_2)^* \subseteq \text{SO}(X, p_1, p_2)$.

Proposition (3-13)

(X, p_1, p_2) is semi- $T_{1/2}$ space if and only if every subset of X is the intersection of all semi-open sets and all semi-closed sets containing it.

Proof

Let (X, p_1, p_2) be a semi- $T_{1/2}$ space with $B \subset X$ arbitrary. then $B = \bigcap \{\{x\}^c : x \notin B\}$ is an intersection of semi-open sets and semi-closed sets by the above theorem the results follow.

Conversely, for each $x \in X$, $\{x\}^c$ is the intersection of all semi-open sets and all semi-closed sets containing it. Thus $\{x\}^c$ is either semi-open sets or semi-closed set and hence X is semi- $T_{1/2}$ space.

References

- 1- J.N.Sharama, "Topology" Meerut college, Meerut.1977.
- 2- J.C.Kelly, "Bitopological spaces" Proc. London Math.Soc. 13 (1963), 71-89.
- 3- K.Balachandran, P.Sundaram and H.Maki, 'on generalized continous maps in topological spaces', Mem.Fac.Sci.Kochi Univ.Ser.A.Math., 12(1991),5-13.
- 4- M. caladas cueva and ratnesh kumar saraf. ' a research on characterizations of semi- $T_{1/2}$ spaces', divulgaciones mtematics vol.1, pp.43-50, 2000
- 5- M. Caldas Cueva, 'semi-generalized continouse maps in topological spaces', portugaliae Mathematical.vol.52 Fasc.4.1995.
- 6- M.K.R.S.Veera Kumar, 'Between semi-closed sets and semi-preclosed sets', J.K.C.college, Guntur-522006, A.P.India, 1991.
- 7- M.Caldas, 'semi- $T_{1/2}$ spaces', Atas sem. Bras.Anal., 40(1994), 191-196.
- 8- M.aldas, 'on g-closed sets and g-continous mappings', Kyungpook Math.jr., 33(2)(1993), 205-209.
- 9- N.Levine, ' semi-open sets and semi-continuity in topological spaces' Amer.math.monthly, 70(1963), 36-41.
- 10- N.Levine, 'generalized closed sets in topology', nd.Circ.Mat.Palermo, 19(1970), 89- 96.
- 11- N.Biswas, 'on characterization of semi-continous function', Atti.Accad.Naz.Lincei Rend.Cl.Sci.Fis.Mat.Natur., 48(8)(1970)399-196.
- 12- P.sundaram, H.Maki, K. Balachandran, 'semi-generalized continous maps and semi- $T_{1/2}$ spaces', Bull.Fukuoka Univ.Ed.part.III, 40(1991), 33-40.
- 13- P.Bhattacharya and B.K.Lahiri, 'semi-generalized closed sets in topology', Indian J.Math., 29(3)(1987), 375-382
- 14- P.Bhattacharyya, and Lahiri, B.K., ' semi-generalized closed sets in topology', Ind.Jr.Math., 29(3)(1987), 375-382.
- 15- R.Devi, H.Maki and K.Balachandran, ' semi-generalized closed maps and generalized semi closed maps', Mem.Fac.Sci.Kochi Univ.Ser.A., 14.(1993), 41-54.

مجلة جامعة كربلاء

مجلة علمية محكمة تعتمد لأغراض الترقيات العلمية تصدر عن رئاسة



رئيس التحرير

أ.د. حسن عودة الغانمي

سكرتير التحرير

أ.م.د. شافي حسين علي الشريفي

مدير التحرير

د. منير عبد العالي عباس الدعيمي

الهيئة الاستشارية	هيئة التحرير
أ.د. حسن عيسى الحكيم	أ.د. عبد عون هاشم الغانمي
أ.د. عباس حسين الحميري	أ.د. حاكم محسن محمد
أ.د. عباس زبون العبودي	أ.م.د. عباس علي محمد الحسيني
أ.د. صاحب علي الاطرقجي	أ.م.د. علاء فراك حسين
أ.د. عبود جودي الحلبي	أ.م.د. علاء الدين الخفاجي
أ.د. عبد عون كاظم السعدي	أ.م.د. علي عبد الحسين الجنابي
أ.عزیز كاظم الناييف	أ.م.د. حسين علي عوض
أ.م.د. عبد الأمير عودة إسماعيل	د. رياض محيي الصائغ
أ.م.د. زهير عمران عيسى	د. أحمد جمعة مهوس
أ.م.د. أحمد عبد الله الراجحي	د. فراس مهدي السلابي

الإدارة الفنية

أسيل عباس حسين إنعام جودة راضي

المشرف اللغوي

محمد عبد الله المهداوي

مجلة جامعة كربلاء العلمية - المجلد السادس - العدد الأول / علمي / ٢٠٠٨