

Republic of Iraq
Ministry of Higher Education
and Scientific Research
University of Babylon
College of Science
Department of Biology



**The Effect of *Pleurotus ostreatus* isolates and AVG on growth
of fenugreek Seeds Under Laboratory Conditions**

A project
Submitted to the Council of the
College of Science University of
Babylon, in Partial Fulfillment of the
Requirements for the Bachelor's
degree in
Biology

By

Fatima Faeiz

Supervised by

Assist. Lecture : Zahraa Abed Neema

بِسْمِ اللّٰهِ الرَّحْمٰنِ الرَّحِیْمِ

يُؤْتِي الْحِكْمَةَ مَنْ يَشَاءُ وَمَنْ يُؤْتَ الْحِكْمَةَ فَقَدْ أُوتِيَ خَيْرًا
كَثِيرًا وَمَا يَذَّكَّرُ إِلَّا أُولُو الْأَلْبَابِ

صدق الله العلي العظيم

آية (269)

Lists of contents

Introduction	4
Method	5
Results	8
discussion	9
References	10

1. Introduction

Fenugreek (*Trigonella foenum graecum L.*) is an annual plant that belongs to the family fabaceae. It is the famous spices in human food. The seeds and green leaves of fenugreek are used in food as well as in medicinal application that is the old practice of human history. It has been used to increase the flavoring and color, and also modifies the texture of food

materials. Fenugreek is known for its pleasantly bitter, slightly sweet seeds. The seeds are available in any form whether whole or ground form is used to flavour many foods mostly curry powders, teas and spice blend. Fenugreek seed has a central hard and yellow embryo which is surrounded by a corneous and comparatively large layer of white and semi-transparent endosperm (Betty, 2008).

The chemical constituents of fenugreek include fibers, saponins, flavonoids, fixed oils, and alkaloids, namely, trigonelline and choline (Shailajan *et al.*, 2011). Additionally, the steroidal sapogenin diosgenin is a secondary metabolite produced by fenugreek and other plants that are widely studied due to its bioactive properties in the medical field (Jesus *et al.*, 2016). The nutritional benefits and curative applications of fenugreek as an effective therapeutic agent against inflammation and diseases, such as cancer and diabetes, have been reported (Syed *et al.*, 2020).

Pleurotus ostreatus (Jacq.) P. Kumm. (*Basidiomycota*), of the *Pleurotaceae* family, comes from China; however, nowadays it is distributed all over the world, except for the north-west Pacific because of the arctic climate. Cultivation methods were developed in Germany during World War I and then successfully applied on a large scale. This was the result of the search for new food sources, due to the problem of hunger in Germany. In Poland, *P. ostreatus* is a common species [Wojewoda 2003].

The efficacy of *Pleurotus ostreatus* has been identified as effective plant-beneficial microbial inoculants, these fungi act as bio-stimulants that promote plant growth (Lombardi *et al.*, 2020). Plant associations with *Pleurotus ostreatus* strains have been shown to enhance plant growth by improving root development, water-holding capacity (Harman *et al.*, 2004), and nutrient

uptake (Yildirim *et al.*, 2006). Also, (Hosseini *et al.*, 2018) , reported that the Fenugreek growth factors and Trigonelline biosynthesis can be affected by *Pleurotus ostreatus* strains. Moreover, it was recently reported that fenugreek inoculation with *Pleurotus ostreatus* treatment increased trigonelline accumulation, which acts as a strong inducing factor for secondary metabolite production (Hosseini *et al.*, 2018).

Aloe leaf extract (ALE) has been used to improve the vegetative growth of *Abelmoschus culentus*, *Oenothera biennis*, and *Majorana hortensis* (Padmaja *et al.*, 2007). Fayeze and Bazaid, (2014)suggested that ALE is an efficient alternative source to improve the growth of *Populus* trees grown under in vitro conditions Paramesha *et al.*, (2021) showed that the early staged fenugreek leaves have the best metabolite and antioxidant capacity. The seeds had higher antioxidant and metabolite activity, but the tender leaves of the Kasuri methi (KS) variety of the plant had the strongest activity for secondary metabolite accumulation.

The aim of this research is Investigate the effect of *Pleurotus ostreatus*, and Aloe vera and there interaction on the growth of Fenugreek plants.

2 . Methods

Source of Fenugreek seeds

Seeds of *T. foenum-graecum L.* were purchased from the local market from Al-Hilla City, Babylon Province, Iraq.

2.1 Preparation Culture media

Potato Dextrose Agar (PDA)

PDA medium was prepared according to the manufacturer's instructions, then sterilized by autoclaved at 121 ° C for 15min. at 15psi and poured in sterilized plates. This medium was used for the growth and maintenance of *Pleurotus ostreatus* isolates (Collee *et al.*, 2015).

2.2 Potato Dextrose Broth (PDB)

PDB medium was prepared by dissolving 24 gm of this medium in 1L of distilled water, mixed thoroughly then serialized and autoclaved.

2.3 Fungal Growth

Pleurotus ostreatus isolates were first cultured on Petri plates containing sterilized PDA, incubated at 26°C for 7 days, in the dark and after that placed under continuous light to promote sporulation

2.4 Sterilization and selection of the fenugreek seeds

Before starting the experiment, all the glassware that we need to prepare treatment formulations were sterilized in an oven for 24 hours at 110 °C. Homogeneous fenugreek seeds, one-year-old with no cracks or other visible deformations were selected, their viability was checked by suspending them in distilled water, then the seeds that settled to the bottom were chosen. Seeds were surface sterilized using 2% NaOCl for 5 min. Seeds were rinsed thoroughly several times with sterile distilled water, the seeds were dried on sterile filter paper under a laminar flow hood. The standard germination of the seeds was 98 %

2.5 Experimental treatment and design

Ten treatments were prepared, in which two isolates of *Pleurotus ostreatus*: Th-1 and Th-2 (1×10^8 spores/ml), and Aloe Vera Gel (AVG) at

50% concentration were conducted. However, ten treatments were applied as the following (the concentration of combination treatments was 50% each):

1. Control, seeds treated with distilled water
2. Seeds treated with (PO-1)
3. Seeds treated with (PO-2)
4. Seeds treated with AVG (100% concentration)
5. Seeds treated with (PO-1) + AVG
6. Seeds treated with (PO-2) + AVG

2.6 Growth parameters

Fenugreek plants were harvested soon after flowering 45 days old, the plant was removed from the pots and the shoots and roots of plants were separated and washed with distilled water three times, and then dried and weighed. and the roots gently washed off from the soil using running tap water. Several leaves, chlorophyll content, shoot length, root length, fresh weight of shoot was evaluated. Shoot length was calculated from the soil line to shoot top with slight modification and then weighed with sensitive balance.

3. Results

The effect of *Pleurotus ostreatus* isolates and AVG on growth of fenugreek seeds under laboratory conditions

Growth parameters of fenugreek plants were varied according to the type of treatment. The results (Table 1) of this experiment revealed that the PO-1 treatment and AVG and their combinations were enhanced the growth capacity of fenugreek plants which summarized as the following:

Root length increased significantly ($P < 0.001$) in plants treated Shoot height, similar tendency was observed with shoot height. Both parts of plant were taller when seeds were treated with PO-1 in lonely.

Fresh weight of shoots was significantly ($P < 0.001$) increased due to seed treatment with PO-1, and AVG and their combinations.

Significant variation was observed in number of leaves in different treatments.

Significant difference was observed in chlorophyll percentage with *Pleurotus ostreatus* treatments.

Table (1). Effects of treatments with *Pleurotus ostreatus* isolates (PO-1 and PO-2) in combination with AVG on root length, shoot height, shoot fresh weight, number of true leaves, and chlorophyll content. of fenugreek (*Pleurotus ostreatus foenum-graecum*) seed treatment Data are means \pm standard errors of four measurements.

Treatments	Root length	Shoot height	FW of of shoots	No. of true leaves	Chlorophyll content
Control	9.12 \pm 0.59	23.30 \pm 1.09	2.10 \pm 0.27	9.20 \pm 0.47	49.07 \pm 0.59
PO-1	14.87 \pm 0.65	32.00 \pm 0.70	3.50 \pm 0.29	12.00 \pm 0.40	56.12 \pm 0.62
PO-2	8.35 \pm 0.69	19.57 \pm 0.33	1.10 \pm 0.29	6.50 \pm 0.86	40.94 \pm 0.33
AVG	14.95 \pm 0.68	31.92 \pm 0.07	7.90 \pm 0.49	23.50 \pm 0.64	59.25 \pm 0.61
PO-1+AVG	14.00 \pm 1.35	33.75 \pm 0.47	6.52 \pm 0.57	28.50 \pm 0.75	61.45 \pm 0.86
PO-2+AVG	10.67 \pm 0.41	20.25 \pm 0.47	2.11 \pm 0.12	8.25 \pm 0.47	45.32 \pm 1.07
L.S.D ₀₀₅	1.3	1.52	0.90	2.38	1.83

4. Discussion

Pleurotus ostreatus showed a wide spectrum of biological activities, among which its hypoglycemic properties are the focus of great interest. *Diabetes mellitus* is one of the most common civilization – related diseases, found both in developed and developing countries. Prognoses indicate an increase in morbidity from year to year [Meetoo et al. 2007].

The present experiment demonstrated the substantial effect of *Pleurotus ostreatus* and its combinations on the growth of fenugreek plants. Since the results of this experiment revealed an increase in root length, shoot height, fresh weight of the shots, number of true leaves, in addition to the chlorophyll content, these parameters reflecting growth improvement (Ezzi & Lynch, 2002).

Some *Pleurotus* species may release soluble compounds and that may exerted beneficial effects not only for plants but also for those microbial groups cohabiting the rhizosphere (Herrera-Jiménez, *et al.*, 2018).

Current experiment revealed that the application of PO-1, and AVG. showed a significant impact on the number of true leaves, shoot length, root length, fresh weight and dry weight of fenugreek plants compared to control. The rapid growth of the plant, number of leaves, plant height, fresh weight and dry weight of plant are the important criterion of growth parameter.

our findings indicated that the effects of PO-1 and AVG alone or in combination with each other on growth mainly depend on the type of *Pleurotus ostreatus* isolate/strain applied. In some cases, *Pleurotus* has a stimulatory effect on plant growth by modifying the soil conditions, besides, the increased growth response of the plant, caused by *Pleurotus ostreatus* strains, mainly depends on the ability to survive and establishment in the rhizosphere (Harman *et al.* 2004). These studies have been confirmed in the case of *Pleurotus ostreatus* enhancing seed germination, root and shoot length (Dubey *et al.* 2007) as well as increasing the frequency of healthy plants. Ozbay and Newman (2004) also reported that *Pleurotus ostreatus* strains have significantly ($P < 0.05$) increased the germination, height, shoot and root dry weight in fenugreek plants transplanted into pots in a greenhouse.).

On other hand, the increasing of plant height in plants that treated with *Pleurotus ostreatus* may due to produce auxins that can stimulate plant growth and root development (Contreras-Cornejo *et al.*, 2009). This result was similar to that obtained by Nagata *et al.*, (2005) who stated that chlorophyll percentage also increased using *Trichoderma* in tomato. For this case, the range of survival percentage was found by *Trichoderma* to improve plant performance under different biotic and abiotic stress (Mastouri 2010).

5. References

- Betty, R.I., 2008. The many healing virtues of fenugreek. *Spice India* 1, 17–19.
- Shailajan, S., Sayed, N., Menon, S., Singh, A., & Mhatre, M. (2011). A validated RP-HPLC method for quantitation of trigonelline from herbal formulations containing *Trigonella foenum-graecum* (L.) seeds. *Pharmaceutical methods*, 2(3), 157-160. doi: 10.4103/2229-4708.90354.
- Syed, Q. A., Rashid, Z., Ahmad, M. H., Shukat, R., Ishaq, A., Muhammad, N., & Rahman, H. U. U. (2020). Nutritional and therapeutic properties of fenugreek (*Trigonella foenum-graecum*): a review. *International Journal of Food Properties*, 23(1), 1777-1791
- Jesus M, Martins A P, Gallardo E. & Silvestre S. (2016). Diosgenin: recent highlights on pharmacology and analytical methodology. *Journal of analytical methods in chemistry*.
- Rajput, S. B., Tonge, M. B., & Karuppayil, S. M. (2014). An overview on traditional uses and pharmacological profile of *Acorus calamus* Linn.(Sweet flag) and other *Acorus* species. *Phytomedicine*, 21(3), 268-276.
- Lombardi, N., Caira, S., Troise, A. D., Scaloni, A., Vitaglione, P., Vinale, F., ... & Woo, S. L. (2020). *Trichoderma* applications on strawberry plants modulate the physiological processes positively affecting fruit production and quality. *Frontiers in microbiology*, 11, 1364.
- Harman, G. E., Doni, F., Khadka, R. B., & Uphoff, N. (2019). Endophytic strains of *Trichoderma* increase plants' photosynthetic capability. *Journal of applied microbiology*.
- Hosseini, Z., Hassanloo, T., Kowsari, M., & Majidian, M. (2018). Trigonelline as an anti-diabetic metabolite increased in inoculated fenugreek by *Trichoderma*. *Advanced Research in Microbial Metabolites & Technology*, 1(2), 129-139.
- Yildirim, E., Taylor, A. G., and Spittler, T. D. (2006). Ameliorative effects of biological treatments on growth of squash plants under salt stress. *Sci. Hortic. (Amst.)* 111:1-6.

- Padmaja, C.K., B.Kowsalya and C. Seethalakshmi (2007) Efficacy of Aloe vera (L.) leaf powder, as a biostimulant in enhancing the growth and yield of lady's finger (*Abelmoschus esculentus* L.). *Research on Crops*,8: 395-397.
- Fayez, K. A., and S. A. Bazaid. 2014. Improving drought and salinity tolerance in barley by application of salicylic acid and potassium nitrate. *Journal of the Saudi Society of Agricultural Sciences* 13 (1):45–55. doi: 10.1016/j.jssas. 2017.01.001.
- Colla, G., Nardi, S., Cardarelli, M., Ertani, A., Lucini, L., Canaguier, R., & Rouphael, Y. (2015). Protein hydrolysates as biostimulants in horticulture. *Scientia Horticulturae*, 196, 28-38.
- Ezzi, M. I. and J. M. Lynch (2002). Cyanide catabolizing enzymes in *Trichoderma* spp. *Enzymol. Microb. Technol.* 31, 1042–1047.
- Rai, N., Limbu, A. K., & Joshi, A. (2020). Impact of *Trichoderma* sp. in Agriculture: A Mini-Review. *Journal of Biology and Today's World*, 9(7), 1-5.
- Herrera-Jiménez, E., Alarcón, A., Larsen, J., Ferrera-Cerrato, R., Cruz-Izquierdo, S., & Ferrera-Rodríguez, M. R. (2018). Comparative effects of two indole-producing *Trichoderma* strains and two exogenous phytohormones on the growth of *Zea mays* L., with or without tryptophan. *Journal of soil science and plant nutrition*, 18(1), 188-201.
- Contreras-Cornejo, H.A.; Macías-Rodríguez, L.; Vergara, A.G.; López-Bucio, J. *Trichoderma* modulates stomatal aperture and leaf transpiration through an abscisic acid-dependent mechanism in *Arabidopsis*. *J. Plant Growth Regul.* **2015**, 34, 425–432. [CrossRef]
- Ozbay, N., & Newman, S. E. (2004). Biological control with *Trichoderma* spp. with emphasis on *T. harzianum*. *Pakistan Journal of Biological Sciences*, 7(4), 478-484.
- Mastouri, F., Björkman, T., & Harman, G. E. (2010). Seed treatment with *Trichoderma harzianum* alleviates biotic, abiotic, and physiological stresses in germinating seeds and seedlings. *Phytopathology*, 100(11), 1213-1221.
- Nagata, N., R. Tanaka, S. Satoh and A. Tanaka. 2005. Identification of a vinyl reductase gene for chlorophyll synthesis in *Arabidopsis thaliana* and

- implications for the evolution of *Prochlorococcus* species. *The Plant Cell*, 17(1): 233-240
- Wojewoda, W. (2003). Checklist of Polish Larger Basidiomycetes. W. Szafer Institute of Botany, Polish Acad. of Sci. Kraków.
- Meetoo, D., McGovern, P., Safadi, R. (2007). An epidemiological overview of diabetes across the world. *Br. J. Nurs.*, 16, 1002–1007.
- Sharma, P.; Jha, A.B.; Dubey, R.S.; Pessarakli, M. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *J. Bot.* **2012**, 2012, 1–26. [CrossRef]
- Marklund .S and Marklund .G, *Eur. J. Biochem.*, 47, 469-474 (1974).
- Frary,A;, Göl, D;, Keleş,D (2010). Salt tolerance in *Solanum pennellii*: antioxidant response and related QTL *BMC Plant Biology* volume 10, Article number: 58 (2010).
- Mir MA, Sirhindi G, Alyemeni MN, Alam P, Ahmad P. Jasmonic acid improves growth performance of soybean under nickel toxicity by regulating nickel uptake, redox balance, and oxidative stress metabolism. *J Plant Growth Regul.* 2018; 37: 1195–1209.