

**Ministry of Higher Education and Scientific Research**

**University of Babylon**

**College of Science**

**Biology department**



LOGO.ADAM96.COM

**A Study the levels of Magnesium, Potassium and Ferritin in  
the Blood Serum of people infected with COVID-19**

**A research submitted to the Department of Biology as part of the  
requirements for obtaining a BSc degree in Biology**

**By**

**Aya kadhim jawad kadhim**

**Supervisor**

**Assist Prof Dr. Zahraa Mohammed Abid Ali Al-Taee**

**2022 A.D**

**1443 A.H**

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

( يَرْفَعِ اللَّهُ الَّذِينَ آمَنُوا مِنْكُمْ وَالَّذِينَ أُوتُوا الْعِلْمَ دَرَجَاتٍ وَاللَّهُ بِمَا

تَعْمَلُونَ خَبِيرٌ ﴿

صدق الله العلي العظيم

[المجادلة: 11]

## Dedication

I dedicate my graduation to the one who harvested thorns from my path, to pave the path of knowledge for me, my dear father, to the one who nursed me with love, tenderness and healing balm, my beloved mother, to the tender hearts and innocent souls of my brothers.

Aya

## Acknowledgement

Praise is to God, who comforted my eyes and my family with my graduation, and thanks are to God who helped me, and on the path of knowledge he patiently endured me.

I would like to extend my thanks and appreciation to the highly ethical Dr. Zahraa Al-Tae for all the valuable guidance and information you have provided us. I ask God to bless and take care of her.

Aya

## *Summary*

---

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the seventh human coronavirus, was discovered in Wuhan, Hubei province, China, during the recent epidemic of pneumonia in January 2020. Since then, the virus has spread all over the world, as of 12 April 2022; there have been 497,960,492 confirmed cases of COVID-19, including 6,181,850 deaths, reported to WHO. SARS-CoV-2 as well as SARS-CoV and Middle East respiratory syndrome coronavirus (MERS-CoV) cause severe pneumonia with a fatality rate of 2.9%, 9.6% and ~36%, respectively.

This study aimed to determine the concentration of some mineral elements (Potassium and Magnesium) and determine the concentration of ferritin.

The current study extended for the period between 1/12/2021 to 1/3/2022, during this period, 89 blood serum samples were collected from people infected with the Corona virus, for ages from 4 years to 64 years, for both sexes (the infection was confirmed by laboratory immunoassay).

It was found that 24% had a decrease in potassium concentration in the blood serum, 11% had an increase in potassium levels, and the remaining 65% had normal potassium levels. This study showed that a number of patients infected with the Corona virus suffer from low levels of magnesium in the blood, and their number was 18 people, including 10 women and two men, while the number of patients suffering from high magnesium in blood serum was 18 infected, including 6 women and 12 men.

## *Summary*

---

It was noted that the percentage of people suffering from a decrease in ferritin protein was 2%, while 3% had a high concentration of this protein in the blood serum, and 94% had a normal level of ferritin.

## List of Constants

Section	Subject	Page No.
	Dedication	I
	Acknowledgment	II
	Summary	III
	List of contents	V
	List of Tables	VI
	List of figures	VII
	List of Abbreviations	VIII
1	Chapter one	1
1.1	Introduction	1
1.2	Aim of study	4
2	Chapter two: Review of literatures	5
2.1	Review of literature	5
2.2	Coronavirus (CoV) structure	6
2.2.1	Symptoms	7
2.2.2	Spread	8
2.2.3	Diagnosis	8
2.3.	Micronutrients	8
2.3.1	Potassium	9
2.3.2	Magnesium (Mg)	10
2.3.3	Iron (Fe)	10
2.3.4	Structure, Regulation and Biological Functions of Ferritin	11
2.3.4.1	Biological Functions of Ferritin	12
3	Chapter three :	13
3.1	Experimental design	13
3.2	Materials	

3.2.1	Devises	14
3.2.2	kit	14
3.3	Methods	15
3.3.1	Samples collection	15
3.3.2	Potassium (Tetraphenylborate Method Without Deproteinization)	15
3.3.3	Magnesium (Xylidyl Blue Monoreagnt)	16
3.3.4	Human Ferritin Heavy Chain, FTH1 ELISA Kit	16
4	Chapter four	21
4.1	Patients with COV-19	21
4.2	Potassium serum concentration	22
4.3	Magnesium serum concentration	25
4.4	Ferritin	26
5	Conclusion and recommendations	29
6	References	30

#### List of Figure

<b>NO. of figure</b>	<b>Title of figure</b>	<b>Page NO.</b>
(2-1)	Remdesivir in the treatment of coronavirus disease 2019 (COVID-19): a simplified summary	7
(2-2)	Ferritin structure: apoferritin forms a roughly spherical container within which ferric iron is stored as a ferrihydrite mineral. Apoferritin refers to the ironfree form of the protein; the iron-containing form is termed holoferritin or simply ferritin. The apoferritin shell is composed of 24 subunits of two types, termed H and L, the ratio of which varies widely depending on tissue type and	12



	inflammation. Iron is toxic in cellular systems because of its capacity to generate reactive species (shown as yellow spheres) which can directly damage DNA and proteins.	
(2-3)	Role of ferritin in different place in human body	14
(3-1)	Dilution of standard solutions	18
(3-2):	A-Before adding stop solution .B- after adding the stop solution	19
(3-3):	Human Ferritin Heavy Chain, FTH1 ELISA Kit	20
(4-1):-	the percentage of male and female to total number	21
(4-2):-	the concentration of potassium in patient with Covid-19.	23
( 4-3 ):-	Compared the concentration of potassium in male and female	23
(4-4):-	the concentration of magnesium in patient with COV-19	25
(4-5):-	the concentration of ferritin protein in patient	27

### List of Tables

NO. of Tables	Title of Tables	Page NO.
(3-1)	Devises used in this study.	14
(3-2)	Kits used in this study.	14
(3-3)	Reaction mixture for serum Potassium concentration	16
(3-4)	Reaction mixture for serum Magnesium concentration	16
(4-1)	Age of patients with Corona virus-19	22

## List of Abbreviations

(IP3)	Inositol 1,4,5-trisphosphate
(DAG)	Diacylglycerol
NK	Natural killer cells
SARS-CoV-2	Severe acute respiratory syndrome coronavirus 2
TMPRSS2	Transmembrane protease, serine 2
CRP	C-reactive protein
ESR	Erythrocyte sedimentation rate
IL-6	InterLeukin-6
TNF-alpha	Tumor necrosis factor
3CLpro	3-chymotrypsin like protease
PLpro	papain like proteases
FTH	Ferritin heavy subunit
FTL	Ferritin light subunit

## *Chapter One*

---

### *Introduction*

Coronavirus disease 2019 (COVID-19) is an infectious disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) Mayo Clinic,2020. It was first identified in December 2019 in Wuhan, China, and has resulted in an ongoing pandemic (Hui DS et al,2020; WHO). The first confirmed case has been traced back to 17 November 2019(Josephine,2020). Traces of the virus have been found in December-2019 wastewater that was collected from Milan and Turin(Kate,2020). As of 21 June 2020, more than 8.8 million cases have been reported across 188 countries and territories, resulting in more than 464,000 deaths. More than 4.37 million people have recovered (Johns Hopkins University,2020).

Minerals are inorganic substances, present in all body tissues and fluids. Their presence is required to maintain certain physical and chemical processes which are essential to life..( Bruulsema *et al*, 2013)

Patients infected with SARS-CoV-2 have been shown to have anomalies in sodium, potassium, chloride, and calcium levels. These alterations in electrolyte balance may be caused by extended hospitalization and the use of various drugs, changes in food habits, multiple organ failure, drug-related adverse effects, and hyperventilation (Alfano *et al*, 2021). There is a strong correlation between electrolyte abnormalities and poor clinical outcomes in Covid19 patients (Ezcan *et al*, 2020; Lippi et al,2020; Sarvazad *et al* ,2020; Wu et al,2020; and arvalho *et al*, 2021).

In confirmed Covid19 instances, potassium problems are the most prevalent electrolyte abnormalities. Both high and low blood potassium concentrations have been discovered to be critical determinants that impact the prognosis of illness independently. Clinical results may be improved if the main cause of potassium homeostasis disturbance is discovered and remedied. (Noori *et al*, 2022). Stabilizing mastocyte membranes in alveolar septa and the submucosal airways

## *Chapter One*

### *Introduction*

---

limit their degranulation, therefore inhibiting their release of many mediators. Toll-like receptor 4/NF- $\kappa$ B axis is hampered by magnesium, which limits neutrophil priming and oxidative burst and also modulates neutrophil and macrophage activity (Maier *et al*, 2020).

Magnesium regulates IP3 and DAG levels in lymphocytes, which are essential second messengers triggered when B and T cell receptors are engaged (Brandao *et al*, 2013).

The cytotoxic activity of T lymphocytes and natural killer (NK) cells is dependent on appropriate intracellular magnesium concentrations, which is why magnesium protects against viral infections (Minton,2013). SARS-CoV-2 infection may be influenced by magnesium levels in the early stages of the disease's progression. Angiotensin-converting enzyme 2 (ACE2), which is expressed in various organs, serves as an entrance receptor for the viral spike (S) protein and is responsible for the pulmonary and extra-pulmonary symptoms of COVID-19 (Gupta *et al*,2020).

Proteolytic cleavage of the S protein by host cell proteases is required for the fusion of viral and cell membranes to occur. Inhibiting the activity of these enzymes may be accomplished in part by the use of magnesium. TMPRSS2 transcription is hampered and, as a result, the enzyme's expression is decreased by magnesium therapy, according to preliminary studies. (Fan *et al*,2021; and Valentina *et al*, 2022).

CRP, ESR, and other inflammatory markers (e.g., IL-6, TNF-alpha, etc.) are elevated in individuals with COVID-19, as are pro-inflammatory cytokines such as IL-6 and TNF-alpha. It has been postulated that the high ferritin level in COVID-19 is connected with the release of cytokines that are involved with inflammation (e.g. IL-6, IL-1 $\beta$ , and TNF-alpha), cellular damage, metabolic acidosis, ROS production, and secondary tissue damage. In individuals with COVID-19, serum ferritin has been found to correlate with disease severity and its surrogates,

## *Chapter One*

---

### *Introduction*

according to individual investigations (CRP) ( Lin *et al* 2020; and Karanvir *et al.*,2022)

#### *Aim of this study:-*

The study aims to determine the relationship of the Corona virus with a number of elements (potassium, magnesium) and to determine the concentration of ferritin during infection with the virus.

## Chapter Two

---

### Review of Literature

#### Review of literature

The current coronavirus disease 2019 (COVID-19) pandemic, caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), which is a member of the Coronavirinae virus subfamily, has exerted an unprecedented global impact. As of a January 13, 2022 update, the world Health Organization COVID-19 Dashboard had reported (312.173) million cases and (5.501) million deaths worldwide (WHO, 2022).

However, when considering “excess deaths” (the standard method of tracking changes in total mortality), 19.2 million (95% confidence interval 12.0–22.4 million) people worldwide are estimated to have died as a consequence of the COVID-19 pandemic (The Economist, 2022). The pandemic catalyzed an extraordinary effort to develop vaccines to induce specific adaptive immunity, based on virus-neutralizing antibodies and T-cell responses. Since their rollout in late 2020, vaccines have been highly effective in preventing clinically significant COVID-19 morbidity/mortality. Not with standing this effect, mutations that have given rise to new variant strains with increased infection transmission rates have emerged to trigger breakthrough infections, even in fully vaccinated individuals (Glatman-Freedman *et al.*,2021; Antonelli *et al.*,2022).

This development has markedly altered the landscape for the management of COVID-19, prompting both renewed “lockdowns” that disrupt local and global economies and the revision of estimates of the level of herd immunity necessary to quell disease.( Kim *et al.*,2021; Niyas *et al.*, 2021) Pressure has also increased on healthcare systems, because of both the reduced efficacy of currently approved vaccines to stop virus transmission, and the chronic disabling symptoms and residual fibrosis in the lungs, kidneys, and cardiovascular system that affect some individuals who have recovered from acute COVID-19 infection—so called “long

## Chapter Two

---

### Review of Literature

COVID.” (Bergwerk *et al.*, 2021; Nalbandian *et al.*, 2021) Consequently, a prudent and, indeed, essential course is to adopt a multipronged strategy that does not place sole reliance on vaccines but instead implements multiple complementary approaches to quell the impact of disease. (Morley and Murray, 2022)

#### Coronavirus (CoV) structure

The CoV family has a sizable homogeneous “spike protein” (Figure 1). The role of the spike (S) protein, which is composed of 1300 amino acids (Elfiky, 2020b), is to interact with the host cells, such as the pulmonary and parabronchial epithelial cell, and assists the coronavirus to enter through the epithelial cell membrane (Boopathi *et al.*, 2020; Xia *et al.*, 2020). Moreover, the alveolar epithelial cells have ample expression of angiotensin-converting enzyme 2 (ACE2), which is an aim by the virus. The detection of ACE2 by the S protein of the virus permits the invasion of the coronavirus into the human circulation system (Belouzard *et al.*, 2009).


Single-strand RNA (22–26 kilobases) viruses such as the coronavirus family reproduce the virus genomes by capitalizing on host cells. For instance, after coronavirus comes near the ribosome of the epithelial cells or other host cells, it utilizes the ribosome of the host cell to replicate poly-proteins. The replication and ensuing procedures of precursor poly-proteins can arise in the epithelial cells (Hoffmann *et al.*, 2020; Wahedi *et al.*, 2020). After the coronavirus polyproteins are exhibited, two enzymes, coronavirus main proteinase (3CLpro) and the papain-like protease (PLpro) are believed to be involved in cleaving the polyproteins into smaller products used for replicating new viruses. In order to produce the daughter RNA genome, the coronavirus exhibits an RNA-dependent RNA polymerase

## Chapter Two

### Review of Literature

(RdRp), which is an important replicase that catalyzes the synthesis of a complementary RNA strand using the virus RNA (Wrapp *et al.*, 2020).

CoVs gather together near intracellular membranes within the Endoplasmic reticulum- Golgi intermediate compartment (or ERGIC) after infection. Here they bud within the lumen and eventually carried outside the cell through ‘exocytosis’ within vesicles (Gupta *et al.*, 2020). Replication of SARS-CoV-2 depends on the viral RNA dependent RNA polymerase (RdRp) (Elfiky & Azzam, 2020) which is the most probable target of the investigational nucleotide analogue **remdesivir (RDV)** (Agostini *et al.*, 2018; Jordan *et al.*, 2018; Siegel *et al.*, 2017; Tchesnokov *et al.*, 2019).



Fig(1-1) Remdesivir in the treatment of coronavirus disease 2019 (COVID-19): a simplified summary

### Symptoms

Common symptoms include fever, cough, fatigue, shortness of breath, and loss of smell and taste (Hopkins, 2020) While the majority of cases result in mild symptoms, some progress to acute respiratory distress syndrome (ARDS) possibly precipitated by cytokine storm (Ye *et al.*, 2020) multi-organ failure, septic shock, and blood clots (Bikdeli *et al.*, 2020) The time from exposure to onset of symptoms



## Chapter Two

---

### Review of Literature

is typically around five days, but may range from two to fourteen days (Velavan and Meyer, 2020)

#### Spread

The virus is primarily spread between people during close contact, most often via small droplets produced by coughing, sneezing, and talking. The droplets usually fall to the ground or onto surfaces rather than travelling through air over long distances. However, research as of June 2020 has shown that speech-generated droplets may remain airborne for tens of minutes. Less commonly, people may become infected by touching a contaminated surface and then touching their face. It is most contagious during the first three days after the onset of symptoms, although spread is possible before symptoms appear, and from people who do not show symptoms (CDC, 2020)

#### Diagnosis

The standard method of diagnosis is by real-time reverse transcription polymerase chain reaction (rRT-PCR) from a nasopharyngeal swab. Chest CT imaging may also be helpful for diagnosis in individuals where there is a high suspicion of infection based on symptoms and risk factors; however, guidelines do not recommend using CT imaging for routine screening (Salehi *et al.*, 2020)

#### Micronutrients

As the world grapples with COVID-19 and the paucity of clinically meaningful therapies, attention has been shifted to modalities that may aid in immune system strengthening.

## Chapter Two

---

### Review of Literature

Taking into consideration that the COVID-19 infection strongly affects the immune system via multiple inflammatory responses, pharmaceutical companies are working to develop targeted drugs and vaccines against SARS- CoV-2 COVID-19.

A balanced nutritional diet may play an essential role in maintaining general wellbeing by controlling chronic infectious diseases. A balanced diet including vitamin A, B, C, D, E, and K, and some micronutrients such as zinc, sodium, potassium, calcium, chloride, and phosphorus may be beneficial in various infectious diseases. A deficiency of vitamins and minerals in the plasma concentration may lead to a reduction in the good performance of the immune system, which is one of the constituents that lead to a poor immune state.

#### Potassium

Hypokalemia can increase ARDS and acute cardiac injury risk, which is considered the most commonly occurring complication in COVID-19. The literature demonstrated that SARS-CoV-2 binds to ACE2 and reduces its expression; consequently, angiotensin-II increases, which subsequently leads to hypokalemia. (Alwaqfi and Ibrahim 2020). COVID- 19 patients showed increased concentration of plasma angi- otensin-II, possibly responsible for acute lung injury and as confirmed in SARS-CoV animal models (Zemlin and Wiese 2020).

A pooled analysis reported that potassium concentration is significantly lower in severe COVID-19 patients than non-severe patients with substantially less heterogeneity than observed for sodium. As with low sodium, reduced plasma potassium levels may be a marker of SARS-CoV-2 infection (Lippi *et al.* 2020).

## Chapter Two

---

### Review of Literature

#### Magnesium (Mg)

Magnesium is the forgotten cation. Mg supplementation might reveal very useful in managing the stress triggered by the pandemic and the post-traumatic stress disorder that will plague COVID-19's survivors, health professionals, and common people. It also plays a significant role in immune function by regulating various functions such as immune cell adherence, immunoglobulin synthesis, binding of Immuno- globulin M (IgM) lymphocyte, antibody-dependent cytotoxicity, and macrophage response towards lymphokines (Ni *et al.* 2020). However, some in vitro and in vivo studies suggest that magnesium plays a vital role in the immune response against viral infections (Jayawardena *et al.* 2020). In Singapore, a cohort study reported that the combination of vitamin D, magnesium, and vitamin B12 (DMB) could reduce the progression rate in older patients with COVID-19. Vitamin B12 (1000 IU) and magnesium (150 mg) have a protective effect against respiratory tract infection and reduce pro-inflammatory cytokines. (Tan *et al.* 2020).

#### Iron (Fe)

Biological function Iron plays a versatile role in the biological system. Despite being an oxidant, iron plays a significant role in hemoglobin and red blood cell production. Role in COVID-19 Recent evidence reveals that apart from pulmonary involvement and elevation in IL-6, COVID- 19 patients display a broader spectrum of hyper-inflammatory syndromes distinguished by cytokine release syndrome (CRS), such as secondary haemophagocytic lymphohistiocytosis (sHLH). Hyperferritinemia is the primary feature of these syndromes,

## Chapter Two

---

### Review of Literature

which plays a significant role in inflammation. These findings support the theory that the acute phase of SARS-CoV-2 infection induces ferritin production associated with the rapid onset of inflammation. Hence, ferritin's immunomodulatory effects contribute to the formation of reactive oxygen species (ROS) and lead to tissue damage. With this contrast, iron chelation therapy is represented as the novel approach against COVID-19. Iron chelation therapy is the most effective approach in a wide spectrum of diseases associated with iron overload. Therefore, iron chelation therapy is considered an appropriate approach to improve survival in COVID-19 patients. A randomized, clinical trial should be considered (da Silva *et al.*,2018).

### **Structure, Regulation and Biological Functions of Ferritin**

Ferritin is a hollow, globular protein of 480 kDa molecular weight. Classical ferritins consist of 24 subunits. Larger forms of ferritins consisting of 36 subunits are present in the heart and skeletal muscles, and microferritin with 12 subunits is produced by bacteria (Andrews *et al.*,2003; Macedo *et al.*, 2003) Mitochondria and nucleus also contain ferritin, which consists of only FTH (da Silva *et al.*,2018)

## Chapter Two

### Review of Literature




Fig. 1-2. Ferritin structure: apoferritin forms a roughly spherical container within which ferric iron is stored as a ferrihydrite mineral. Apoferritin refers to the iron-free form of the protein; the iron-containing form is termed holoferritin or simply ferritin. The apoferritin shell is composed of 24 subunits of two types, termed H and L, the ratio of which varies widely depending on tissue type and inflammation. Iron is toxic in cellular systems because of its capacity to generate reactive species (shown as yellow spheres) which can directly damage DNA and proteins.

### Biological Functions of Ferritin

The main function of ferritin is to store iron (Tosha *et al.*, 2012). FTH has ferroxidase activity, which converts soluble ferrous iron into storable ferric iron. Then, ferric iron enters the cavity of ferritin and forms an iron core under the action of the nucleation site of FTL. Ferritin rich in FTH can accumulate and release iron faster than that rich in FTL, and has a more active iron transport system. Ferritin rich in FTL has more iron than that rich in FTH, which plays an important role in iron storage (Hasan *et al.*, 2008). Christophe Ferreira examined the embryonic

## Chapter Two

---

### Review of Literature

lethality of mice that knocked out the FTH gene, and the results showed that FTH<sup>-/-</sup> embryos died at 3.5–9.5 days of embryonic development (Ferreira *et al.* 2000). And FTH silencing disrupts the formation of S-S bonds, causing miss folding of the protein, and increase in the cellular ROS level (Zolea *et al.*,2015). However, FTL<sup>-/-</sup> mice showed systemic and brain iron dyshomeostasis (Li *et al.*, 2019). This indicated that the expression of the FTH could compensate the function of the FTL to sequester iron in vivo. In addition to iron storage, ferritin functions as a protective agent preventing iron overload, decreasing oxidative stress. It has been reported that NF- $\kappa$ B antagonizes the tumor necrosis factor- $\alpha$  (TNF- $\alpha$ )-induced apoptosis by upregulating the expression of FTH. The protective effect of ferritin is achieved through iron sequestration and suppressing ROS accumulation, thus inhibiting C-Jun N-terminal kinase (JNK) signaling (Antosiewicz *et al.*,2007); And ferritin is also reported as an acute phase protein, which is part of a general pattern of the systemic inflammation (Slaats *et al.*,2016 ; Moreira *et al.* 2020).

In the early stage of inflammation, pro-inflammatory cytokines such as interleukin-1 (IL-1) stimulates the expression and secretion of ferritin, which reduces the level of iron and inhibits the apoptosis induced by IL-1 (Festa *et al.* ,2015). Studies have shown that the function of ferritin is more dynamic than previous depicted. Ferritin also acts as an immuno-suppressor. Alterations of H-ferritin/T cell immunoglobulin and mucin domain-containingprotein-2 (Tim-2) binding/signaling might be involved in thepathogenesis of autoimmune diseases. Therefore, the detection of ferritin concentration is not only an important indicator to diagnose diseases with iron overload or iron deficiency, but also a marker of inflammatory conditions or autoimmune disorders (Recalcati *et al.*, 2008).

# Chapter Two

## Review of Literature





Fig (1-3) Role of ferritin in different place in human body

*Chapter Three*  
*Material and Methods*

---

*Experimental design*





**Chapter Three**  
**Material and Methods**

❖ Materials :-

A- Devises :-

Devises used in this study listed in table (3-1)

Table (3-1) Devises used in this study.

No.	Apparatus	Company	Country
.1	centrifuge	Hettich	Germany
.2	Different size Micropipettes $\mu$ l	Dragon	German
.3	Different size tips	Biobaseic	Canada
.4	Different size tube	Biobaseic	Canada
.5	ELISA reader & washer	Biotek_USA	USA
.6	Gel tube		
.7	Spectrophotometer		
.8	Syringe	Meheco	China

B- Kit

Kits used in this study listed in table (3-2)

Table (3-2) Kits used in this study.

No.	KIT	CATALOG NO.	COMPANY
1.	Potassium (Tetraphenylborate Method Without	298 002	spectrum

## Chapter Three

### Material and Methods

	Deproteinization)		
2.	MAGNESIUM (Xylidyl Blue Monoreagnt)	285 002	spectrum
3.	Human Ferritin Heavy Chain, FTH1 ELISA Kit	E6766Hu	BT LAB

#### ❖ Methods

##### 1- Samples Collection:-

Blood samples (5ml) by sterile medical syringe were collected from people infected with Corona virus, their ages ranged between 4-64 years and of both sexes. All volunteers were subjected to a number of questions as indicated in the information form in the appendix. The samples were kept in special gel tubes to separate the blood serum. The tubes were left at room temperature for half an hour, after which they were centrifuged with a centrifuge device to separate the blood serum and stored at a temperature  $-20^{\circ}\text{C}$ .

##### 2- Potassium (Tetraphenylborate Method Without Deproteinization)

In order to measure the potassium concentration in the blood serum, the following steps were taken:

- a- Dissolve the serum in a water bath at  $30^{\circ}\text{C}$ .
- b- The reaction mixture was prepared according to the table (3-3)

## Chapter Three

### Material and Methods

Table (3-3) Reaction mixture for serum Potassium concentration

	Reagent Blank	Standard	Sample
Reagent R	1mL	1mL	1mL
Standard	.....	20µl	.....
Sample	.....	.....	20µl

- c- The mixtures were mixed, and incubated for 3 minutes at 37 °C.
- d- Mixed again thoroughly and read absorbance of sample ( $A_{\text{sample}}$ ) and standard ( $A_{\text{standard}}$ ) against blank on 578nm.
- e- The following equation was used to find the potassium concentration in the blood serum (samples)

$$\text{Sreum Potassium Conc. (mmol/L)} = \frac{A_{\text{sample}}}{A_{\text{standard}}} \times 5$$

### 3- Magnesium (Xylidyl Blue Monoreagnt) :-

The following steps were taken to measure the concentration of magnesium in the blood serum:

- a- Dissolve the serum in a water bath at 30°C.
- b- The reaction mixture was prepared according to table (3-4)

Table (3-3) Reaction mixture for serum Potassium concentration

	Blank	Standard	Sample
Reagent R	1mL	1mL	1mL
Standard	.....	10µl	.....
Sample	.....	.....	10µl

## *Chapter Three*

---

### *Material and Methods*

- c- All mixtures were mixed well and let to stand for 10 minutes at room temperature, then absorbance (at 546nm) of sample and standard read against reagent blank.
- d- The following equation was used to find the magnesium concentration in the blood serum (samples)


$$\text{serum Magnesium Conc. (mg/dl)} = \frac{A \text{ sample}}{A \text{ standard}} \times 2.5$$

#### **4- Human Ferritin Heavy Chain, FTH1 ELISA Kit**

- a- Reagents, standard solutions and samples were prepared according to the following steps.
  - i. All reagents were thawed and brought to room temperature before use
  - ii. Standard was Reconstitute the 120ul of the standard (320ng/ml) with 120ul of standard diluent to generate a 160ng/ml standard stock solution. Allow the standard to sit for 15 min. with gentle agitation prior to making dilutions. Prepare duplicate standard points by serially diluting the standard stock solution (160ng/ml) 1:2 with standard diluent to produce 80ng/ml, 40ng/ml, 20ng/ml and 10ng/ml solutions. Standard diluent serves as the zero standards (0ng/ml). Dilution of standard solutions figure (3-1)

**Chapter Three**  
**Material and Methods**

---



**Figure (3-1) Dilution of standard solutions**

iii. **Wash Buffer** Dilute 20ml of Wash Buffer Concentrate 25x into deionized or distilled water to yield 500 ml of 1x Wash Buffer. If crystals have formed in the concentrate, mix gently until the crystals have completely dissolved.

iv. **Samples preparation**

- Samples were within 5 days (stored at 2-8°C).
- Samples were brought to room temperature before starting the assay.
- Centrifuge to collect sample before use.
- The supernatants were collected carefully.

Before using any reagents, bring them to room temperature. Assays are completed in a room-temperature environment.


b- A 50ul of standard was added to standard well.

c- A 40ul of sample was added to sample wells and then 10ul of Human FTH1 antibody added to sample wells, then 50ul streptavidin-HRP added to sample

## Chapter Three

### Material and Methods

- wells and standard wells. The Mixtures mixed well, the plate covered with a sealer and incubate for 60 minutes at 37°C.
- d- the sealer Remove and the plate washed 5 times with wash buffer. wells Soaked with 300ul wash buffer for 30 seconds to 1 minute for each wash. the plate dried onto paper towels or other absorbent material.
  - e- A 50ul of substrate solution A added to each well and then 50ul of substrate solution B added to each well. plate covered with a new sealer and Incubated for 10 minutes at 37°C in the dark.
  - f- A 50ul of Stop Solution added to each well, the blue color would change into yellow immediately fig1.



**Figure (3-2): A-Before adding stop solution .B- after adding the stop solution**

**Chapter Three**  
**Material and Methods**

---

g- The optical density (OD value) of each well Determined immediately using a microplate reader set to 450 nm within 10 minutes after adding the stop solution.




**Figure (3-3): Human Ferritin Heavy Chain, FTH1 ELISA Kit**

**1- Patients with COV-19:-**

An 89 blood serum samples were taken from people infected with Corona virus from both women and men, for the purpose of studying the concentration of the number of elements (potassium, magnesium and ferritin) that are expected to be affected as a result of infection with the virus.

The number of women was 35(20%) and the number of men was 54 (30%), age range was (4-64 years) as in Figure 1



**Figure (1):- the percentage of male and female to total number**

The subjects included in this study were divided into the following groups according to age as shown in Table (1). A questionnaire (included age, duration, number of infection with COV-19, treatment and other disease) was designed to obtain the information of patient's subjects (depending on AUA score). The laboratory work was carried out in the research laboratory in the Department of Biology, College of Science, University of Babylon.



Table (1): Age of patients with Corona virus-19

Age	No. of cases
1-----15	8
16-----30	24
31-----45	28
46 ----- 60	13
61 ----- 75	17

**2- Potassium serum concentration:-**

Among the 89 samples studied in this research, it was found that 24% had a decrease in potassium concentration in the blood serum, 11% had an increase in potassium levels, and the remaining 65% had normal potassium concentration , Fig. 2. When comparing the potassium concentration between male and female, it was noted that the increase in potassium was more in male than in female. As for the low potassium concentration, it was more in female than in male than in male Fig.3.

Potassium deficiency is common among Covid-19 patients because this element is excreted in large quantities in the urine, so most patients suffer from potassium deficiency.

Chen *et al.*, 2020, came to the conclusion that people with COVID-19 tend to have hypokalemia. Hypokalemia is hard to fix because the kidneys lose K<sup>+</sup> all the time because ACE2 (which is a SARS-CoV-2 receptor) is broken down. The powerful vasoconstrictor angiotensin II is turned into angiotensin by the enzyme Angiotensin-Converting Enzyme 2. Because of this action, ACE2 is involved in

## Chapter Four

### Result and Discussion

the renin-angiotensin system. The end of urine  $K^+$  loss is a good sign for the patient's prognosis and may be a reliable, in-time, and sensitive biomarker that shows the end of the bad effect on the RAS system.




Figure (2):- the concentration of potassium in patient with Covid-19.




Figure (3):- Compared the concentration of potassium in male and female

## *Chapter Four*

### *Result and Discussion*

---

Hypokalemia happened more often than usual in COVID-19 patients who were very sick. A recent Chinese study found that 93 percent of COVID-19 patients had low potassium levels (Chen et al,2020). When ACE2 is turned down in people with COVID-19, it causes a very bad imbalance in the renin-angiotensin system. As biomarkers of hypokalemia, people who were out of balance lost less potassium in their urine and responded more to intravenous potassium (Silhol, et al, 2020).

When people with severe hypokalemia were given 3 g of potassium per day (an average of 34 g over the course of a hospital stay), they got better. Hypokalemia, which is when the potassium level in the blood is below 3.5 mEq/L, can cause problems with fluid and electrolytes in the clinic. 50% of people with clinically confirmed low magnesium levels also had low magnesium levels (Whang et al., 1992). However, the serum level used to define deficiency varies from 0.6 to 0.75 mmol/L, with some being more cautious than others (Huang and Kuo ,2007 ; and Veltri and Mason,2015). Hypomagnesemia can aggravate hypokalemia and induce renal potassium loss. When you don't get enough magnesium, the sodium-potassium ATPase pump doesn't work as well. This lowers the amount of potassium inside the cell. When there isn't enough magnesium in the body, distal potassium release through ROMK channels makes more potassium waste. In magnesium deficiency, increased salt transport to the extremities and aldosterone levels may make potassium loss worse. However, this is not the only thing that can cause hypokalemia (Huang and Kuo ,2007).

Certain medications (e.g., antibiotics, beta 2-receptor antagonists, diuretics, insulin, glucocorticoids, laxatives, etc) may also cause hypokalemia through a variety of mechanisms including intracellular potassium shifting, increased renal loss, and decreased intestinal uptake. Deficiency in magnesium in critically ill patients can not only lead to secondary hypokalemia but also hypocalcemia,


## Chapter Four

### Result and Discussion

accompanied by severe neuromuscular and cardiovascular clinical manifestations. Concurrent magnesium deficiency aggravates hypokalemia and renders it refractory to potassium treatment (Veltri and Mason,2015). If the patient is hypomagnesemic or has borderline low-normal serum concentrations, treating hypomagnesemia concomitantly with the hypokalemia is appropriate to prevent serum potassium levels from falling below the normal range after discontinuation of potassium therapy (Wallace,2020).

#### **Magnesium serum concentration:-**

This study showed that a number of patients infected with the Corona virus suffer from low levels of magnesium in the blood, and their number was 18 people, including 10 women and two men, while the number of patients suffering from high magnesium in blood serum was 18 infected, including 6 women and 12 men Fig(4).



**Figure (4):- the concentration of magnesium in patient with COV-19**

Among the many essential nutrients, magnesium is the second most abundant intracellular cation after potassium, and involved in >600 enzymatic reactions in

## *Chapter Four*

### *Result and Discussion*

---

the body, including those contributing to the exaggerated immune and inflammatory responses exhibited by COVID-19 patients (Wallace,2020).


Magnesium deficiency makes phagocytes more active, raises granulocyte oxidative stress, activates endothelial cells, and raises cytokine levels, all of which lead to inflammation (Wallace *et al.*,2020). Several parts of the COVID-19 are similar to the metabolic changes that have been shown to happen during latent subclinical magnesium deficiency. These changes include a drop in T cells, an increase in the amount of inflammatory cytokines in the blood, and problems with the endothelium. Our data showed that severe COVID-19 patients had significantly lower magnesium levels than non-severe COVID-19 patients. This is in line with the idea by Iotti *et al.*,2021 that a low Mg status might contribute to the change from mild to severe COVID-19 symptoms. It was thought that keeping an eye on the amount of magnesium in the body could be a way to slow or stop the spread of disease (Hao-Long *et al.*, 2021).

#### **Ferritin:-**

When measuring the level of ferritin in the blood serum, it was noted that the percentage of people suffering from a decrease in this protein was 2%, while 3% had a high level of ferritin in the blood serum, and 94% had a normal level of ferritin. Figure 5

## Chapter Four

### Result and Discussion



**Figure (5):- the concentration of ferritin protein in patient**

Most of the research indicates an increase in ferritin concentration in the blood serum when infected with the Corona virus.

Following the global spread of the COVID-19 pandemic and early throughout the pandemic, elevated levels of ferritin in patients with COVID-19 brought the attention of medical staff treating huge numbers of patients (Wang *et al* 2020) as well as researchers by predicting poor outcomes (Huang *et al.*, 2020). The possible correlation between ferritin and worse prognosis however, barely scratched the surface of the hidden elements beneath it which necessitated further investigation. Subsequently, ferritin concentration progressed from a prognostic point of view, to serve as a severity risk factor. More importantly, the dramatic increase in ferritin concentration, alongside lymphopenia, reduced NK cell number activity, abnormal liver function tests, and coagulopathy, had researchers speculating, and later agreeing that COVID-19 might be the newest member in the group of hyperferritinemic syndromes (Colafrancesco *et al.*,2020 and Naim *et al.*, 2022)

## *Chapter Four*

---

### *Result and Discussion*

Patients with severe to critical disease had higher ferritin concentration than those with mild to moderate disease, and non-survivors had higher serum ferritin concentration than survivors. Patients requiring ICU and mechanical ventilation had higher serum ferritin concentration than those who didn't require ICU and didn't require mechanical ventilation, according to Kaushal *et al*,2022. COVID-19 patients with renal involvement had a higher concentration of ferritin as well; however, there was heterogeneous data about the relationship between ferritin concentration and COVID-19-associated liver damage. A decreased blood ferritin concentration was seen in patients who did not have COVID-19-related thrombotic events. Serum ferritin is an important indicator of severity in COVID-19 (Kaushal *et al*,2022).

## Conclusions and Recommendations

---

### ***Conclusion:-***

- 1- The number of people infected with the virus with low potassium level was more than patients with high potassium concentration.
- 2- More women than men suffer from low levels of magnesium.
- 3- Within the population studied, it was found that patients infected with coronavirus who had an elevated ferritin protein concentration were slightly more than those with a deficiency of this protein.

### ***Recommendations:-***

Studying the effect of different vaccines for the Corona virus on minerals and micronutrients in the human body.



## References

---

- "Coronavirus disease 2019 (COVID-19)—Symptoms and causes". Mayo Clinic .Retrieved 14 April 2020.
- "COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU)". ArcGIS.Johns Hopkins University.Retrieved 21 June 2020.
- "WHO Director-General's opening remarks at the media briefing on COVID- 19" World Health Organization (WHO) (Press release). 11 March 2020. Archived from the original on 11 March 2020.Retrieved 12 March 2020.
- Agostini, M. L., Andres, E. L., Sims, A. C., Graham, R. L., Sheahan, T. P., Lu, X., Smith, E. C., Case, J. B., Feng, J. Y., Jordan, R., Ray, A. S., Cihlar, T., Siegel, D., Mackman, R. L., Clarke, M. O., Baric, R. S., & Denison, M. R. (2018). Coronavirus susceptibility to the antiviral remdesivir (GS-5734) is mediated by the viral polymerase and the proofreading exoribonuclease. *mBio*, 9(2), 1–15. <https://doi.org/10.1128/mBio.00221-18>.
- Alfano G, Ferrari A, Fontana F, et al. Hypokalemia in patients with COVID-19. *Clin Exp Nephrol*. 2021;25:1-409. <https://doi.org/10.1007/s10157-020-01996-4>
- Alwaqfi NR, Ibrahim KS. COVID-19: an update and cardiac involvement. *J Cardiothorac Surg*. 2020 Sep 9;15(1):239. doi: 10.1186/s13019-020-01299-5. PMID: 32907637; PMCID: PMC7480628.
- Andrews SC, Robinson AK, Rodríguez-Quñones F (2003) Bacterial iron homeostasis. *FEMS Microbiol Rev* 27(2-3):215–237. [https://doi.org/10.1016/s0168-6445\(03\)00055-x](https://doi.org/10.1016/s0168-6445(03)00055-x)
- Antonelli M, Penfold RS, Merino J, et al. Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom

## References

---

- Study. app: a prospective, community-based, nested, case control study. *Lancet Infect Dis.* 2022;22:43–55.
- Antosiewicz J, Ziolkowski W, Kaczor JJ, Herman-Antosiewicz A (2007) Tumor necrosis factor-alpha-induced reactive oxygen species formation is mediated by JNK1-dependent ferritin degradation and elevation of labile iron pool. *Free Radic Biol Med* 43(2):265–270. <https://doi.org/10.1016/j.freeradbiomed.2007.04.023>
- Belouzard, S., Chu, V. C., & Whittaker, G. R. (2009). Activation of the SARS
- Bergwerk M, Gonen T, Lustig Y, et al. Covid-19 breakthrough infections in vaccinated health care workers. *N Engl J Med.* 2021;385:1474–1484.
- Bikdeli B, Madhavan MV, Jimenez D, Chuich T, Dreyfus I, Driggin E, et al. (April 2020). "COVID-19 and Thrombotic or Thromboembolic Disease: Implications for Prevention, Antithrombotic Therapy, and Follow-up". *Journal of the American College of Cardiology.*75 (23): 2950–2973. doi:10.1016/j.jacc.2020.04.031. PMC 7164881.PMID 32311448.
- Boopathi, S., Poma, A. B., & Kolandaivel, P. (2020). Novel 2019 coronavirus
- Brandao K, Deason-Towne F, Perraud A-L, Schmitz C (2013) The role of Mg<sup>2+</sup> in immune cells. *Immunol Res* 55:261–269. <https://doi.org/10.1007/s12026-012-8371-x>
- Bruulsema, T., P. Heffer, R. Welch, I. Cakmak, and K. Moran. 2012. Fertilizing Crops to Improve Human Health: A Scientific Review. *Better Crops with Plant Food* 96(2):29-31.
- Chen D, Li X, Qifa s, Hu C, Su F, Dai J. Hypokalemia and clinical implications in patients with coronavirus disease 2019 (COVID-19). *Infectious Diseases (except HIV/AIDS).* 2020. doi: 10.1101/2020.02.27.20028530

## References

---

Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Xia J, Yu T, Zhang X, Zhang L. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020 Feb 15;395(10223):507-513. doi: 10.1016/S0140-6736(20)30211-7. Epub 2020 Jan 30. PMID: 32007143; PMCID: PMC7135076.

coronavirus spike protein via sequential proteolytic cleavage at two distinct sites. *Proceedings of the National Academy of Sciences of the United States of America*, 106(14), 5871–5876. <https://doi.org/10.1073/pnas.0809524106>.

D. Wang, B. Hu, C. Hu, F. Zhu, X. Liu, J. Zhang, et al., Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in wuhan, China, *J. Am. Med. Assoc.* 323 (11) (2020) 1061–1069

da Silva VK, de Freitas BS, Dornelles VC, Kist LW, Bogo MR, Silva MC, Streck EL, Hallak JE et al (2018) Novel insights into mitochondrial molecular targets of iron-induced neurodegeneration: reversal by cannabidiol. *Brain Res Bull* 139:1–8. <https://doi.org/10.1016/j.brainresbull.2018.01.014>

De Carvalho H, Richard MC, Chouihed T, et al. Electrolyte imbalance in COVID-19 patients admitted to the Emergency Department: a case-control study. *Intern Emerg Med*. 2021:1-6. <https://doi.org/10.1007/s11739-021-02632-z14>.

Elfiky, A. A. (2020b). SARS-CoV-2 RNA dependent RNA polymerase (RdRp) targeting: An in silico perspective. *Journal of Biomolecular Structure and Dynamics*, 1–15. <https://doi.org/10.1080/07391102.2020.1761882>

Elfiky, A. A., & Azzam, E. B. (2020). Novel guanosine derivatives against MERS CoV polymerase: An in silico perspective. *Journal of Biomolecular*

## References

---

- Structure and Dynamics, 1–9. <https://doi.org/10.1080/07391102.2020.1758789>.
- Ezcan ME, DoganGokce G, Sen N, Zorlutuna Kaymak N, Ozer RS. Baseline electrolyte abnormalities would be related to poor prognosis in hospitalizedcoronavirusedisease2019patients.NewMicrobesNew Infect.2020;37:100753.<https://doi.org/10.1016/j.nmni.2020.10075311>.
- Fan L, Zhu X, Zheng Y, et al. (2021) Magnesium treatment on methylation changes of transmembrane serine protease 2(TMPRSS2). medRxiv 03.11.21253287
- Ferreira C, Bucchini D, Martin ME, Levi S, Arosio P, Grandchamp B, Beaumont C (2000) Early embryonic lethality of H ferritin gene deletion in mice. J Biol Chem 275(5):3021–3024. <https://doi.org/10.1074/jbc.275.5.3021>
- Festa L, Gutoskey CJ, Graziano A, Waterhouse BD, Meucci O (2015) Induction of interleukin-1 beta by human immunodeficiency virus-1 viral proteins leads to increased levels of neuronal ferritin heavy chain, synaptic injury, and deficits in flexible attention. J Neurosci 35(29):10550–10561. <https://doi.org/10.1523/Jneurosci.4403-14.2015>
- Giuseppe Lippi, Fabian Sanchis-Gomar, Brandon M. Henry.2020. Coronavirus disease 2019 (COVID-19): the portrait of a perfect storm. Annals of Translational Medicine 8(7)
- Glatman-Freedman A, Bromberg M, Dichtiar R, et al. The BNT162b2 vaccine effectiveness against new COVID-19 cases and complications of breakthrough cases: a nation-wide retrospective longitudinal multiple cohort analysis using individualised data. EBioMedicine. 2021;72:103574.

## References

---

- Gordon, C. J., Tchesnokov, E. P., Woolner, E., Perry, J. K., Feng, J. Y., Porter, D. P., & Gotte, M. (2020). Remdesivir is a direct-acting antiviral that inhibits RNA-dependent RNA polymerase from severe acute respiratory syndrome coronavirus 2 with high potency. *Journal of Biological Chemistry*, 295(15), 4773–4779. <https://doi.org/10.1074/jbc.RA120.013679>.
- Gupta A, Madhavan MV, Sehgal K et al (2020) Extrapulmonary manifestations of COVID-19. *Nat Med* 26:1017–1032. <https://doi.org/10.1038/s41591-020-0968-3>
- Gupta, M. K., Vemula, S., Donde, R., Gouda, G., Behera, L., & Vadde, R. (2020). In-silico approaches to detect inhibitors of the human severe acute respiratory syndrome coronavirus envelope protein ion channel. *Journal of Biomolecular Structure and Dynamics*, 1–11. <https://doi.org/10.1080/07391102.2020.1751300>.
- Hao-Long Zeng, Qing Yang, Peihong Yuan, Xu Wang and Liming Cheng. 2020. Associations of essential and toxic metals/metalloids in whole blood with both disease severity and mortality in patients with COVID-19. *The FASEB Journal*. ;35. <https://doi.org/10.1096/fj.202002346RR>
- Hasan MR, Tosha T, Theil EC (2008) Ferritin contains less iron (Fe-59) in cells when the protein pores are unfolded by mutation. *J Biol Chem* 283(46):31394–31400. <https://doi.org/10.1074/jbc.M806025200>
- Hoffmann, M., Kleine-Weber, H., Krüger, N., Müller, M., Drosten, C., & Hopkins C. "Loss of sense of smell as marker of COVID-19 infection". Ear, Nose and Throat surgery body of United Kingdom. Retrieved 28 March 2020
- Huang C-L, Kuo E. Mechanism of hypokalemia in magnesium deficiency. *J Am Soc Nephrol*. 2007;18(10):2649–52. doi:10.1681/ASN.2007070792.

## References

---

- Huang, R. Pranata, M.A. Lim, A. Oehadian, B. Alisjahbana, C-reactive protein, procalcitonin, D-dimer, and ferritin in severe coronavirus disease-2019: a meta-analysis, *Ther. Adv. Respir. Dis.* 14 (2020), 1753466620937175.
- Hui DS, I Azhar E, Madani TA, Ntoumi F, Kock R, Dar O, et al. (February 2020). "The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health—The latest 2019 novel coronavirus outbreak in Wuhan, China". *International Journal of Infectious Diseases*.91: 264–266. doi:10.1016/j.ijid.2020.01.009. PMC 7128332.PMID1953166.
- Iotti S, Wolf F, Mazur A, Maier JA. The COVID-19 pandemic: is there a role for magnesium? hypotheses and perspectives. *Magnes Res.* 2020; 33(2):21–27. <https://doi.org/10.1684/mrh.2020.0465>
- Jayawardena R, Sooriyaarachchi P, Chourdakis M, Jeewandara C, Ranasinghe P (2020) Enhancing immunity in viral infections, with special emphasis on COVID-19: A review. *Diabetes MetabSyndr Clin Res Rev.* <https://doi.org/10.1016/j.dsx.2020.04.015>
- Karanvir Kaushal, , Hardeep Kaur, et al.(2022). Serum ferritin as a predictive biomarker in COVID-19. A systematic review, meta-analysis and meta-regression analysis. *Journal of Critical Care* 67 :172–181 . <https://doi.org/10.1016/j.jcrc.2021.09.023>
- Kelland, Kate (19 June 2020). "Italy sewage study suggests COVID-19 was there in December 2019". REUTERS. Retrieved 20 June 2020.
- Kim JH, Hotez P, Batista C, et al. Operation Warp Speed: implications for global vaccine security. *Lancet Global Health.* 2021;9:e1017–e1021.

## References

---

- Li W, Garringer HJ, Goodwin CB, Richine B, Acton A, VanDuyn N, Muhoberac BB, Irimia-Dominguez J et al (2015) Systemic and cerebral iron homeostasis in ferritin knock-out mice. *PLoS One* 10(1):ARTN e0117435. <https://doi.org/10.1371/journal.pone.0117435>
- Lin Z, Long F, Yang Y, Chen X, Xu L, Yang M. Serum ferritin as an independent risk factor for severity in COVID-19 patients. *J Infect.* 2020;81:647–79. <https://doi.org/10.1016/j.jinf.2020.06.053>.
- Lippi G, SouthAM, HenryBM. Electrolyte imbalances in patients with severe coronavirus disease 2019 (COVID-19). *Ann Clin Biochem.* 2020;57:262-265. <https://doi.org/10.1177/000456322092225512>.
- Lu, L. (2020). Fusion mechanism of 2019-nCoV and fusion inhibitors targeting HR1 domain in spike protein. *Cellular and Molecular Immunology*, 1–3.
- Ma, Josephine (13 March 2020). "Coronavirus: China's first confirmed Covid-19 case traced back to November 17". *South China Morning Post*. Archived from the original on 13 March 2020. Retrieved 28 May 2020.
- Macedo S, Romão CV, Mitchell E, Matias PM, Liu MY, Xavier AV, LeGall J, Teixeira M et al (2003) The nature of the di-iron site in the bacterioferritin from *Desulfovibrio desulfuricans*. *Nat Struct Biol* 10(4):285–290. <https://doi.org/10.1038/nsb909>
- Maier JA, Castiglioni S, Locatelli L et al (2020) Magnesium and inflammation: advances and perspectives. *Semin Cell Dev Biol* S1084–9521(20):30171–30173. <https://doi.org/10.1016/j.semdb.2020.11.002>
- Maier JA, Castiglioni S, Locatelli L, Zocchi M, Mazur A. Magnesium and inflammation: advances and perspectives. *Semin*

## References

---

- Cell Dev Biol. 2020;S1084-9521(20):30171–30173. <https://doi.org/10.1016/j.semcd.2020.11.002>
- Maryam Noori, Seyed A. Nejadghaderi, Mark J. M. Sullman, Kristin Carson-Chahhoud, Ali-Asghar Kolahi and Saeid Safiri. Epidemiology, prognosis and management of potassium disorders in Covid-19. *Rev Med Virol.*,2022. DOI: 10.1002/rmv.2262
- Minton K (2013) Immunodeficiency: magnesium regulates antiviral immunity. *Nat Rev Immunol* 13:548–549. <https://doi.org/10.1038/nri3501>
- Moreira AC, Mesquita G, Gomes MS (2020) Ferritin: an inflammatory player keeping iron at the core of pathogen-host interactions. *Microorganisms* 8(4):ARTN 589. <https://doi.org/10.3390/microorganisms8040589>
- Morley D. Hollenberg and Murray Epstein. 2022. The innate immune response, microenvironment proteinases, and the COVID-19 pandemic: pathophysiologic mechanisms and emerging therapeutic targets. *Kidney International Supplements* ,12, 48–62; <https://doi.org/10.1016/j.kisu.2021.12.001>
- Mukhtar H. Ahmed<sup>1</sup> · Arez Hassan<sup>2</sup> · Judit Molnár.2021. The Role of Micronutrients to Support Immunity for COVID-19 Prevention. *Revista Brasileira de Farmacognosia* 31:361–3. <https://doi.org/10.1007/s43450-021-00179-w>
- Naim Mahroum, Amal Alghory, Zeynep Kiyak, Abdulkarim Alwani, Ravend Seida, Mahmoud Alrais, Yehuda Shoenfeld, 2022. Ferritin – from iron, through inflammation and autoimmunity, to COVID-19, *Journal of Autoimmunity*,vol.126. <https://doi.org/10.1016/j.jaut.2021.102778>.



## References

---

- Nalbandian A, Sehgal K, Gupta A, et al. Post-acute COVID-19 syndrome. *Nat Med.* 2021;27:601–615.
- Ni W, Yang X, Yang D, Bao J, Li R, Xiao Y et al (2020) Role of angiotensin-converting enzyme 2 (ACE2) in COVID-19. *Crit Care* 24(1):1–10. <https://doi.org/10.1186/s13054-020-03120-0>
- Niyas VKM, Arjun R. Correspondence: breakthrough COVID-19 infections among health care workers after two doses of ChAdOx1 nCoV-19 vaccine. *QJM.* 2021;114:757–758.
- Pöhlmann, S. (2020). The novel coronavirus 2019 (2019-nCoV) uses the SARS-coronavirus receptor ACE2 and the cellular protease TMPRSS2 for entry into target cells. *BioRxiv*, 1–23. <https://doi.org/10.1101/2020.01.31.929042>
- Recalcati S, Invernizzi P, Arosio P, Cairo G (2008) New functions for an iron storage protein: the role of ferritin in immunity and autoimmunity. *J Autoimmun* 30(1-2):84–89. <https://doi.org/10.1016/j.jaut.2007.11.003>
- S. Colafrancesco, C. Alessandri, F. Conti, R. Priori, COVID-19 gone bad: a new character in the spectrum of the hyperferritinemic syndrome? *Autoimmun. Rev.* 19 (7) (2020) 102573
- Salehi S, Abedi A, Balakrishnan S, Gholamrezanezhad A (March 2020). "Coronavirus Disease 2019 (COVID-19): A Systematic Review of Imaging Findings in 919 Patients". *AJR. American Journal of Roentgenology*: 1–7. doi:10.2214/AJR.20.23034. PMID 32174129.
- SarvazadH, CahngaripourSH, EskandariRoobahaniN, Izadi B.Evaluation of electrolyte status of sodium, potassium and magnesium, and

## References

---

- fasting blood sugar at the initial admission of individuals with COVID-19 without underlying disease in Golestan Hospital Kermanshah. *New Microbes New Infect.* 2020;38:100807. <https://doi.org/10.1016/j.nmni.2020.100807>
- Siegel, D., Hui, H. C., Doerffler, E., Clarke, M. O., Chun, K., Zhang, L., Neville, S., Carra, E., Lew, W., Ross, B., Wang, Q., Wolfe, L., Jordan, .Soloveva, V., Knox, J., Perry, J., Perron, M., Stray, K. M., Barauskas, O., ... Mackman, R. L. (2017). Discovery and synthesis of a phosphoramidate prodrug of a pyrrolo[2,1-f][triazin-4-amino] adenine C-nucleoside (GS-5734) for the treatment of Ebola and emerging viruses. *Journal of Medicinal Chemistry*, 60(5), 1648–1661. <https://doi.org/10.1021/acs.jmedchem.6b01594>.
- Silhol F, Sarlon G, Deharo J-C, Vaïsse B. Downregulation of ACE2 induces overstimulation of the renin–angiotensin system in COVID-19: should we block the renin–angiotensin system? *Hypertens Res.* 2020. doi:10.1038/s41440-020-0476-3.
- Slaats J, ten Oever J, van de Veerdonk FL, Netea MG (2016) IL-1 beta/IL-6/CRP and IL-18/ferritin: distinct inflammatory programs in infections. *PLoS Pathog* 12(12):ARTN e1005973. <https://doi.org/10.1371/journal.ppat.1005973>
- Stadnytskyi V, Bax CE, Bax A, Anfinrud P (June 2020). "The airborne lifetime of small speech droplets and their potential importance in SARS-CoV-2 transmission". *Proceedings of the National Academy of Sciences of the United States of America*. 117 (22): 11875–11877. doi:10.1073/pnas.2006874117. PMC 7275719. PMID 32404416.

## References

---

- structure, mechanism of action, antiviral drug promises and rule out against its treatment. *Journal of Biomolecular Structure and Dynamics*, 1–14. <https://doi.org/10.1080/07391102.2020.1758788>
- Tan CW, Ho LP, Kalimuddin S, Cherng BPZ, Teh YE, Thien SY et al (2020) A cohort study to evaluate the effect of combination Vitamin D, Magnesium and Vitamin B12 on progression to severe outcome in older COVID-19 patients. *Nutrition*. <https://doi.org/10.1101/2020.06.01.20112334>
- Tchesnokov, E. P., Feng, J. Y., Porter, D. P., & Gotte, M. (2019). Mechanism of inhibition of Ebola virus RNA-dependent RNA polymerase by remdesivir. *Viruses*, 11(4), 326. <https://doi.org/10.3390/v11040326>
- Tosha T, Behera RK, Theil EC (2012) Ferritin ion channel disorder inhibits Fe(II)/O-2 reactivity at distant sites. *Inorg Chem* 51(21):11406–11411. <https://doi.org/10.1021/ic3010135>
- Valentina Trapani· Andrea Rosanoff· Shadi Baniyadi · et al(2022). The relevance of magnesium homeostasis in COVID-19. *European Journal of Nutrition*. 61:625–636 <https://doi.org/10.1007/s00394-021-02704-y>
- Velavan TP, Meyer CG (March 2020). "The COVID-19 epidemic". *Tropical Medicine & International Health*. 25 (3): 278–280. doi:10.1111/tmi.13383. PMC 7169770.PMID 32052514.
- Veltri KT, Mason C. Medication-induced hypokalemia. *P T Peer Rev J Formul Manag*. 2015;40(3):185–90.
- Wahedi, H. M., Ahmad, S., & Abbasi, S. W. (2020). Stilbene-based natural compounds as promising drug candidates against COVID-19. *Journal of*

## References

---

- Biomolecular Structure and Dynamics, 1–16. <https://doi.org/10.1080/07391102.2020.1762743>.
- Wallace TC. Combating COVID-19 and Building Immune Resilience: A Potential Role for Magnesium Nutrition? *J Am Coll Nutr.* 2020 Nov-Dec;39(8):685-693. doi: 10.1080/07315724.2020.1785971. Epub 2020 Jul 10. PMID: 32649272.
- Whang R, Whang DD, Ryan MP. Refractory potassium repletion. A consequence of magnesium deficiency. *Arch Intern Med.* 1992;152(1):40–5.
- World Health Organization. WHO Coronavirus (COVID-19) Dashboard. Accessed January 13, 2022. <https://covid19.who.int/>
- Wrapp, D., Wang, N., Corbett, K. S., Goldsmith, J. A., Hsieh, C.-L., Abiona, O., Graham, B. S., & McLellan, J. S. (2020). Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science (New York, N.Y.)*, 367(6483), 1260–1263. <https://doi.org/10.1126/science.abb2507>.
- Wu Y, Hou B, Liu J, Chen Y, Zhong P. Risk factors associated with long-term hospitalization in patients with COVID-19:a single-centered, retrospective study. *Front Med.* 2020;7.<https://doi.org/10.3389/fmed.2020.00315>
- Xia, S., Zhu, Y., Liu, M., Lan, Q., Xu, W., Wu, Y., Ying, T., Liu, S., Shi, Z., Jiang, S., &
- Ye Q, Wang B, Mao J (June 2020). "The pathogenesis and treatment of the 'Cytokine Storm' in COVID-19". *J. Infect.* 80 (6): 607–613. doi:10.1016/j.jinf.2020.03.037. PMC 7194613.PMID 32283152
- Zemlin AE, Wiese OJ. Coronavirus disease 2019 (COVID-19) and the renin-angiotensin system: A closer look at angiotensin-converting enzyme 2 (ACE2). *Ann Clin Biochem.* 2020 Sep;57(5):339-350. doi:

## References

---

- 10.1177/0004563220928361. Epub 2020 Jun 2. PMID: 32369402; PMCID: PMC7267743.
- Zhang, N., Yu, X., Xie, J. *et al.* (2021). New Insights into the Role of Ferritin in Iron Homeostasis and Neurodegenerative Diseases. *Mol Neurobiol* 58, 2812–2823 <https://doi.org/10.1007/s12035-020-02277-7>.
- Zolea F, Biamonte F, Candeloro P, Di Sanzo M, Cozzi A, Di Vito A, Quaresima B, Lobello N et al (2015) H ferritin silencing induces protein misfolding in K562 cells: a Raman analysis. *Free Radic Biol Med* 89:614–623. <https://doi.org/10.1016/j.freeradbiomed.2015.07.161>
- Centers for Disease Control and Prevention (CDC). "How COVID-19 Spreads". 2 April 2020. Archived from the original on 3 April 2020. Retrieved 3 April 2020.
- Salehi S, Abedi A, Balakrishnan S, Gholamrezanezhad A (March 2020). "Coronavirus Disease 2019 (COVID-19): A Systematic Review of Imaging Findings in 919 Patients". *AJR. American Journal of Roentgenology*: 1–7. doi:10.2214/AJR.20.23034. PMID 32174129 .

استمارة معلومات

رقم العينة:-

تأريخ جمع العينة:-

الملاحظات	الاسم
	العمر
	الجنس
	مدخن
	يتناول الكحول
	هل يعاني من ارتفاع ضغط الدم
	هل يعاني من مرض السكري
	هل يعاني من امراض اخرى (تذكر)
	عدد المرات التي اصيب بها الشخص بـ COVID

## الخلاصة

تم اكتشاف فيروس كورونا 2 (SARS-CoV-2) ، الفيروس التاجي البشري السابع ، في ووهان بمقاطعة هوبي بالصين ، خلال وباء الالتهاب الرئوي الأخير في يناير 2020. ومنذ ذلك الحين انتشر الفيروس في جميع أنحاء العالم اعتبارًا من 12 أبريل 2022 ، كان هناك 492,960,497 حالة مؤكدة من COVID-19 ، بما في ذلك 850,6181 حالة وفاة وتم إبلاغ منظمة الصحة العالمية بها. يتسبب فيروس SARS-CoV-2 وكذلك فيروس SARS-CoV ومتلازمة الشرق الأوسط التنفسية (MERS-CoV) في حدوث التهاب رئوي حاد بمعدل وفيات يبلغ 2.9% و 9.6% و 36% على التوالي.

هدفت هذه الدراسة لتحديد تركيز بعض العناصر المعدنية (البوتاسيوم والمغنيسيوم والفيريتين) في الجسم ودراسة وجود رابط بينها وبين الإصابة بالفايروس.

امتدت الدراسة الحالية للفترة بين 2021/12/1 الى 2022/3/1 حيث تم خلال هذه الفترة جمع ٨٩ عينة مصل الدم من الاشخاص المصابين بفايروس كورونا للاعمار من ٤ سنوات الى ٦٤ سنة ولكلا الجنسين.

وجد أن 24% لديهم انخفاض في تركيز البوتاسيوم في مصل الدم ، 11% لديهم زيادة في مستويات البوتاسيوم ، و 65% المتبقية لديهم مستويات طبيعية من البوتاسيوم. أظهرت هذه الدراسة أن عددا من المرضى المصابين بفايروس كورونا يعانون من انخفاض مستويات المغنيسيوم في الدم ، وكان عددهم 18 شخصا بينهم 10 نساء ورجلين ، فيما كان عدد المرضى الذين يعانون من ارتفاع المغنيسيوم في مصل الدم 18 مصابا بينهم 6 نساء و 12 رجلا. كما تم قياس مستوى الفيريتين لدى الاشخاص المصابين حيث لوحظ أن نسبة الأشخاص الذين يعانون من نقص هذا البروتين كانت 2% ، بينما 3% لديهم مستوى عالٍ من الفيريتين في مصل الدم ، و 94% لديهم مستوى طبيعي من الفيريتين.



LOGO.ADAM96.COM

وزارة التعليم العالي والبحث العلمي

جامعة بابل

كلية العلوم

قسم علوم الحياة

دراسة مستويات كل من المغنيسيوم والبوتاسيوم والفرتين في مصل الدم

لاشخاص مصابين بكوفيد-19

بحث مقدم الى قسم علوم الحياة كجزء من متطلبات نيل شهادة البكلوريوس تخصص علوم الحياة  
بواسطة

آيه كاظم جواد كاظم

باشراف

أ.م.د. زهراء محمد عبد علي الطائي